
GENERAL TOPOLOGY

A COLLECTION OF NOTES ON MAJOR DEFINITIONS AND RESULTS, PROOFS, AND
COMMENTARY BASED ON THE CORRESPONDING COURSE AT ILLINOIS, AS INSTRUCTED BY
MINEYEV

LECTURE NOTES BY
DHEERAN E. WIGGINS

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Disclaimer

The lecture notes in this document were based on General Topology [535], as instructed by Igor Mineyev [Department of Mathematics] in the Fall semester of 2024 [FA24] at the University of Illinois Urbana-Champaign. All non-textbook approaches, exercises, and comments are adapted from Mineyev's lectures.

Textbook

Many of the exercises and presentations were selected from *Topology, Second Edition*, by James R. Munkres.

Author Information

Dheeran E. Wiggins is, at the time of writing [Fall, 2024], a second-year student at Illinois studying mathematics. All typesetting and verbiage are his own.

All mathematics begins with the empty set.

– Igor Mineyev

Contents

Contents	v
ON TOPOLOGY, CONTINUITY, AND INVARIANTS	1
1 Topological Preliminaries	3
1.1 Notations	3
1.2 Building a Plane Topology	4
1.3 Spaces and Bases	5
1.4 Comparing Topologies	6
1.5 New Topologies From Old	7
1.6 Homeomorphisms and Continuity	9
1.7 Product Topology	11
1.8 Closure, Interior, and Boundary	13
1.9 Aside on Norms and Sequences of Functions	15
2 Connectedness and Compactness	21
2.1 (Path-)Connectedness	21
2.2 Compactness	24
2.3 Local and Sequential Compactness	26
2.4 Lebesgue's Number Lemma	28
2.5 Compactifications and Zorn's Lemma	30
ON EMBEDDINGS, FUNCTION SPACES, AND ALGEBRAIC TOPOLOGY	35
3 Separation and Embeddings	37
3.1 Introducing the T_i Axioms	37
3.2 Embeddings and Stone-Čech	38
4 Completeness and Function Spaces	41
4.1 Cauchy Sequences and Completions	41
4.2 Bounded Space	42
4.3 More on Function Spaces	45
4.4 Arzelà-Ascoli Theorem	48
5 Introduction to Algebraic Topology	49
5.1 Fundamental Group	50
5.2 Covering Spaces	53

ON TOPOLOGY, CONTINUITY, AND INVARIANTS

Topological Preliminaries

1

1.1 Notations

- 1.1 Notations 3
- 1.2 Building a Plane Topology . 4
- 1.3 Spaces and Bases 5
- 1.4 Comparing Topologies 6
- 1.5 New Topologies From Old . 7
- 1.6 Homeomorphisms and
Continuity 9
- 1.7 Product Topology 11
- 1.8 Closure, Interior, and
Boundary 13
- 1.9 Aside on Norms and
Sequences of Functions . . . 15

For the purpose of this course, the set of real numbers \mathbb{R} will look like the real line, not paying as close attention to the precise logical model. Similarly, $\mathbb{R}^2 := \mathbb{R} \times \mathbb{R}$, $\mathbb{R}^3 := \mathbb{R} \times \mathbb{R} \times \mathbb{R}$, and so forth.

Definition 1.1.1 (2-Sphere) *We define the set*

$$\mathbb{S}^2 := \{x \in \mathbb{R}^3 : d(0, x) = 1\},$$

taking the standard Euclidean metric.

Following suit, we can define the n -sphere.

Definition 1.1.2 (n -Sphere)

$$\mathbb{S}^n := \{x \in \mathbb{R}^{n+1} : d(0, x) = 1\}.$$

Now, you will notice that our n -sphere lives in $n + 1$ -space. In what sense is \mathbb{S}^2 two-dimensional? Well, after doing some groundwork, we can show that *locally*, the 2-sphere is akin to a plane.

Remark 1.1.1 Generally, we will write a set as X , and the corresponding power set as $\mathcal{P}(X) := \{S : S \subseteq X\}$.

Definition 1.1.3 (Family) *A family of subsets of X is any subset $\mathcal{F} \subseteq \mathcal{P}(X)$.*

Definition 1.1.4 (Family Intersection) *The intersection of a family $\mathcal{F} \subseteq \mathcal{P}(X)$ is¹*

$$\bigcap_{S \in \mathcal{F}} S := \{x : \text{for all } S \in \mathcal{F}, x \in S\}.$$

1: The union is defined similarly.

Definition 1.1.5 (Open Ball) *An open ball $B(x, r)$ of radius $r > 0$ around $x \in E$ is defined as²*

$$B(x, r) := \{y \in E : d(x, y) < r\}.$$

2: We take a metric space (E, d) .

1.2 Building a Plane Topology

Returning to the plane \mathbb{R}^2 , let us look at the families

$$\mathcal{B} := \{B(x, r) : x \in \mathbb{R}^2, r \in (0, \infty)\} \subseteq \mathcal{P}(\mathbb{R}^2)$$

$$\mathcal{T}(\mathcal{B}) := \left\{ \bigcup \mathcal{B}' : \mathcal{B}' \subseteq \mathcal{B} \right\} \subseteq \mathcal{P}(\mathbb{R}^2).$$

Clearly $\mathcal{B} \subseteq \mathcal{T}$, because for all $B(x, r)$, $\mathcal{T} \ni \bigcup \{B(x, r)\} = \{B(x, r)\}$. Now, consider $(0, 1) \times (0, 1) \subset \mathbb{R}^2$.³ Now, $B(x, r_x) \subseteq (0, 1) \times (0, 1)$, so

$$(0, 1) \times (0, 1) = \bigcup_{x \in (0,1) \times (0,1)} B(x, r_x) \in \mathcal{T}.$$

In general, we can find such open balls for any “similar” shape. We will develop the notion of open sets by identifying them with elements of \mathcal{T} .

3: This is simply the open unit square. We can fill the open square with open balls via $B(x, r_x)$, where a formula for r_x comes from the infimum of the distance between x and a point on the boundary.

4: That is, \mathcal{T} is closed under arbitrary unions.

5: Take $\mathcal{B}_U \subseteq \mathcal{B}$.

6: Is this possible? Well, we can pick the radius

$$r_x := \min \left\{ \begin{array}{l} r_1 - d(x_1, x) \\ r_2 - d(x_2, x) \end{array} \right\}$$

where $B_1 = B(x_1, r_1)$ and $B_2 = B(x_2, r_2)$. Proving that $B(x, r_x) \subseteq B_1 \cap B_2$ is just an exercise in the triangle inequality.

7: In this case, $\overline{B}(0, 1)$ is the closed ball, not the closure of the open ball.

Proposition 1.2.1

- (i) For all $U_1, U_2 \in \mathcal{T}$, $U_1 \cup U_2 \in \mathcal{T}$.
- (ii) For any subfamily $\mathcal{F} \subseteq \mathcal{T}$, $\bigcup \mathcal{F} \in \mathcal{T}$.⁴
- (iii) For all $U_1, U_2 \in \mathcal{T}$, $U_1 \cap U_2 \in \mathcal{T}$.
- (iv) For any finite subfamily of $\mathcal{F} \subseteq \mathcal{T}$, $\bigcap \mathcal{F} \in \mathcal{T}$.

(i) *Proof.* We know $U_1 = \bigcup \mathcal{B}_1$ and $U_2 = \bigcup \mathcal{B}_2$ for some families $\mathcal{B}_1, \mathcal{B}_2$. Then,

$$U_1 \cup U_2 = \left(\bigcup \mathcal{B}_1 \right) \cup \left(\bigcup \mathcal{B}_2 \right) = \bigcup (\mathcal{B}_1 \cup \mathcal{B}_2) \in \mathcal{T}.$$

□

(ii) *Proof.* Take any $\mathcal{F} \subseteq \mathcal{T}$. Now, $\bigcup \mathcal{F} = \bigcup_{U \in \mathcal{F}} U$. We know U is a union of balls in \mathbb{R}^2 , so $U = \bigcup \mathcal{B}_U$.⁵ Then,

$$\bigcup \mathcal{F} = \bigcup_{U \in \mathcal{F}} U = \bigcup_{U \in \mathcal{F}} \left(\bigcup \mathcal{B}_U \right) = \bigcup_{B \in \bigcup_{U \in \mathcal{F}} \mathcal{B}_U} B = \bigcup \mathcal{F}' \in \mathcal{T}.$$

□

(iii) *Proof.* We have $U_1 = \bigcup \mathcal{B}_1$ and $U_2 = \bigcup \mathcal{B}_2$, as before, so

$$U_1 \cap U_2 = \left(\bigcup \mathcal{B}_1 \right) \cap \left(\bigcup \mathcal{B}_2 \right) = \bigcup_{\substack{B_1 \in \mathcal{B}_1 \\ B_2 \in \mathcal{B}_2}} (B_1 \cap B_2).$$

For each pair B_1, B_2 , choose a family \mathcal{F}_{B_1, B_2} of open balls such that $\bigcup \mathcal{F}_{B_1, B_2} = B_1 \cap B_2$. To construct \mathcal{F}_{B_1, B_2} , for all $x \in B_1 \cap B_2$, choose a sufficiently small radius r_x such that $B(x, r_x) \subseteq B_1 \cap B_2$.⁶ Then, $B_1 \cap B_2 = \bigcup_{x \in B_1 \cap B_2} B(x, r_x)$, so $U_1 \cap U_2 \in \mathcal{T}$, given our knowledge about unions.

(iv) Use induction. □

Remark 1.2.1 We cannot prove that for *any* subfamily, their intersection lies in \mathcal{T} . For instance, consider $B_\varepsilon := B(0, 1 + \varepsilon)$ and $\mathcal{F} := \{B_\varepsilon : \varepsilon > 0\}$. Well, $\bigcap \mathcal{F} = \overline{B}(0, 1) \notin \mathcal{T}$.⁷

We also trivially have that $\emptyset \in \mathcal{T}$ and $\mathbb{R}^2 \in \mathcal{T}$, taking unions as needed.⁸

8: These properties will provide motivation for the definition of a topology \mathcal{T} .

1.3 Spaces and Bases

Definition 1.3.1 (Metric Space) Recall that a metric space is an ordered pair (X, d) , where X is a set and $d : X \times X \rightarrow [0, \infty)$ such that⁹

- (i) $d(x, y) = 0$ if and only if $x = y$.
- (ii) $d(x, y) = d(y, x)$.
- (iii) $d(x, y) + d(y, z) \geq d(x, z)$.

9: The function d is called the *metric* or *distance* function. The third property is the *triangle inequality*.

Topologies can loosen our conditions so that we no longer need a metric function d on the set. Topologies are, thus, *weaker* structures than metrics.

Definition 1.3.2 (Topology) A topology on a set X is any family \mathcal{T} of subsets of X such that,

- (i) $\emptyset, X \in \mathcal{T}$
- (ii) \mathcal{T} is closed under arbitrary unions.
- (iii) \mathcal{T} is closed under finite intersections.

Remark 1.3.1 Every metric induces a topology, following the form of the proof from the discussion of the plane topology.¹⁰

10: We form our topology by unioning the subfamilies of all possible balls.

Definition 1.3.3 (Topological Space) The ordered pair (X, \mathcal{T}) is called a topological space.¹¹

11: We refer to the set X as the *underlying* set of the topological space.

Remark 1.3.2 As is usual with ordered tuple structures, we will prefer to write a space (X, \mathcal{T}) as simply X .¹²

12: This is only valid when the topology is clear for context, or clarified at the beginning of a statement.

Definition 1.3.4 (Open Set) A subset $U \subseteq X$ is called an open set if $U \in \mathcal{T}$.

Definition 1.3.5 (Closed Set) A subset $C \subseteq X$ is called closed if $X \setminus C \in \mathcal{T}$.¹³

13: That is, a set is closed if its complement is open.

Definition 1.3.6 (Basis of Topology) Let X be a set. A basis \mathcal{B} of a topology on X is a family of subsets¹⁴ such that

- (i) for all $x \in X$, there exists $B \in \mathcal{B}$, such that $x \in B$.¹⁵
- (ii) for all $B_1, B_2 \in \mathcal{B}$, for any $x \in B_1 \cap B_2$, there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.¹⁶

14: Remember, this means $\mathcal{B} \subseteq \mathcal{P}(X)$.

15: Equivalently, $\bigcup \mathcal{B} = X$.

16: That is, \mathcal{B} should be “dense enough.” Equivalently, for any $B_1, B_2 \in \mathcal{B}$, there exists a subfamily $\mathcal{B}' \subseteq \mathcal{B}$ such that $\bigcup \mathcal{B}' = B_1 \cap B_2$.

Based on our proof in \mathbb{R}^2 , it is clear to see that the metric gives a basis, which provides us with a topology. In general, we simply need the basis to generate a topology.

17: To prove both topologies are equal, simply prove both directions of inclusion.

Definition 1.3.7 (Topology Generated by a Basis) *Let X be a set and \mathcal{B} be a basis of a topology. The following are equivalent definitions of the topology generated by the basis $\mathcal{T}(\mathcal{B})$:*¹⁷

(i)

$$\mathcal{T}(\mathcal{B}) := \{U \subseteq X : \text{for all } x \in U \text{ there is } B \in \mathcal{B} \text{ s.t. } x \in B \subseteq U\}.$$

(ii)

$$\mathcal{T}(\mathcal{B}) := \left\{ \bigcup \mathcal{B}' : \mathcal{B}' \subseteq \mathcal{B} \right\}.$$

Proposition 1.3.1 *The generated topology $\mathcal{T}(\mathcal{B})$ is a topology.*

Proof. Follow the recipe shown for the plane. □

1.4 Comparing Topologies

Example 1.4.1 Let $X := [0, 1] \subset \mathbb{R}$. We could define a topology

$$\mathcal{T}_1 := \{\emptyset, [0, 1]\} \subset \mathcal{P}([0, 1]).$$

18: In general, the trivial topology on any set X is just $\mathcal{T} := \{\emptyset, X\}$. It is the smallest, via cardinality, possible topology. Similarly, the discrete topology on X is $\mathcal{T} := \mathcal{P}(X)$

This is the *trivial* topology on $[0, 1]$.¹⁸ We could also define another topology $\mathcal{T}_2 := \mathcal{P}([0, 1])$. This is called the *discrete* topology on $[0, 1]$.

Now that we have multiple topologies, it is natural to wish to compare them. Let \mathcal{T}_1 and \mathcal{T}_2 be topologies on X .

Definition 1.4.1 (Finer/Stronger Topology) *If we have $\mathcal{T}_1 \subseteq \mathcal{T}_2$, then we say \mathcal{T}_1 is smaller/courser/weaker and \mathcal{T}_2 is larger/finer/stronger.*

Remark 1.4.1 An equivalent way to characterize the discrete topology is to take a basis defined by

$$\mathcal{B}_{\text{disc}} := \{\{x\} : x \in X\},$$

the set of singletons.¹⁹ Then, $\mathcal{T}(\mathcal{B}_{\text{disc}})$ is the discrete topology on X .

19: Sometimes you may see this called the *discrete* basis.

Lemma 1.4.1 (Basis Criterion) *Let \mathcal{T} be a topology on X and let $\mathcal{C} \subseteq \mathcal{T}$. Then, the following are equivalent:*

- (i) *For any $U \in \mathcal{T}$ and for all $x \in U$, there exists $B \in \mathcal{C}$ such that $x \in B \subseteq U$.*
- (ii) *\mathcal{C} is a basis, and $\mathcal{T}(\mathcal{C}) = \mathcal{T}$.*

Proof. For (i) \Rightarrow (ii), take any $x \in X$. We have that $X \in \mathcal{T}$, so there exists a $B \in \mathcal{C}$ so that $x \in B \subseteq X$. Thus, $\bigcup \mathcal{C} = X$. Take any $B_1, B_2 \in \mathcal{C}$. Suppose $x \in B_1 \cap B_2$. Since \mathcal{T} is a topology, $B_1 \cap B_2 \in \mathcal{T}$. Thus, there

exists $B_3 \in \mathcal{C}$ such that $x \in B_3 \subseteq U$, so \mathcal{C} is a basis. It remains to be shown that $\mathcal{T}(\mathcal{C}) = \mathcal{T}$, but this is easy. We leave (ii) \Rightarrow (i) as an exercise. \square

1.5 New Topologies From Old

Definition 1.5.1 (Subspace Topology) *Let (X, \mathcal{T}) be a topological space. Suppose $A \subseteq X$. We define the subspace topology*

$$\mathcal{T}_A := \{A \cap U : U \in \mathcal{T}\} \subseteq \mathcal{P}(A).$$

Then, (A, \mathcal{T}_A) is the induced subspace of the ambient space (X, \mathcal{T}) .

Proposition 1.5.1 \mathcal{T}_A , as above, is a topology.²⁰

Proof. We have that $\emptyset \in \mathcal{T}_A$, as $\emptyset = A \cap \emptyset$. Similarly, $A \in \mathcal{T}_A$, as $A = A \cap X$. Pick any subfamily of \mathcal{T}_A . It must be of the form $\{A \cap U : U \in \mathcal{F}\}$, for some $\mathcal{F} \subseteq \mathcal{T}$. Well,

$$\bigcup_{U \in \mathcal{F}} (A \cap U) = A \cap \bigcup_{U \in \mathcal{F}} U = A \cap \bigcup \mathcal{F} \in \mathcal{T}_A.$$

Finally, taking \mathcal{F} to be finite this time,

$$\bigcap_{U \in \mathcal{F}} (A \cap U) = A \cap \bigcap_{U \in \mathcal{F}} U = A \cap \bigcap \mathcal{F} \in \mathcal{T}_A.$$

\square

Lemma 1.5.2 *Let (X, \mathcal{T}) be a topological space, and let $A \subseteq X$. If $\mathcal{T} = \mathcal{T}(\mathcal{B})$ for some basis \mathcal{B} in X , then consider $\mathcal{B}_A := \{A \cap B : B \in \mathcal{B}\}$. Then, \mathcal{B}_A is a basis for \mathcal{T}_A .²¹*

Proof. The proof is left as an exercise. \square

We now consider a nice canonical construction.

Definition 1.5.2 (Set Disjoint Union) *Given two sets X, Y , the disjoint union $X \coprod Y$ is the union of a bijective copy of X and bijective copy of Y , such that the copies are disjoint.²²*

One way of creating these copies is to take $\{0, 1\} \times X \cup Y$, where we only keep X in the 0 slice and Y in the 1 slice.

Lemma 1.5.3 *The family $\mathcal{T}_X \cup \mathcal{T}_Y$ is a basis of a topology on $X \coprod Y$.*

Definition 1.5.3 (Disjoint Union Topology) *Let $\{(X_i, \mathcal{T}_i)\}_{i \in \Lambda}$ be an*

20: This is a very powerful construction, giving us a way to form as many topological subspaces as we like.

21: Remember, this is *two* statements. We have that \mathcal{B}_A is a basis of a topology, and that $\mathcal{T}(\mathcal{B}_A) = \mathcal{T}_A$.

22: However, you will also see the disjoint union used for when $X \cap Y = \emptyset$.

arbitrary family of topological spaces. Then, the topology on

$$X := \coprod_{i \in \Lambda} X_i$$

is precisely the topology \mathcal{T} generated by the union:

$$\mathcal{T} := \mathcal{T} \left(\bigcup_{i \in \Lambda} \mathcal{T}_i \right).$$

23: That is, for all $x \in X$, there is an $S \in \mathcal{S}$ such that $x \in S$.

Definition 1.5.4 (Subbasis) *A subbasis on a set X is a family $\mathcal{S} \subseteq \mathcal{P}(X)$ such that $\bigcup \mathcal{S} = X$.²³*

Remark 1.5.1 Any basis is a subbasis.

24: We also need S' to be nonempty.

Definition 1.5.5 (Basis From Subbasis) *Given a subbasis \mathcal{S} on X , we can produce another family $\mathcal{B}(\mathcal{S})$ defined by²⁴*

$$\mathcal{B}(\mathcal{S}) := \left\{ \bigcap \mathcal{S}' : \mathcal{S}' \subseteq \mathcal{S} \text{ and } \mathcal{S}' \text{ is finite} \right\}.$$

We can now form a topology generated by the subbasis $\mathcal{T}(\mathcal{B}(\mathcal{S}))$.

25: Say f is surjective.

Definition 1.5.6 (Topology Induced by Function) *Let (X, \mathcal{T}_X) be a topological space, and let $f : X \rightarrow Y$, where Y is just a set.²⁵ Define an induced topology*

$$\mathcal{T}_Y := \{V \subseteq Y : f^{-1}(V) \in \mathcal{T}_X\}.$$

26: Convince yourself that there is a homeomorphism between the circle and $[0, 1]/\sim$.

Example 1.5.1 Put any equivalence relation \sim on $[0, 1]$. Identify $0 \sim 1$. Then, $[0, 1]/\sim$ can be visualized as a circle. We can construct a map $q : [0, 1] \rightarrow [0, 1]/\sim$ such that $x \mapsto [x]$. Clearly, q is surjective.²⁶

Definition 1.5.7 (Quotient Topology) *Given a space (X, \mathcal{T}_X) and an equivalence relation \sim on X . Then, the quotient topology on X/\sim is the topology induced by the canonical quotient map. That is,*

$$\mathcal{T}_{X/\sim} := \{V \subseteq X/\sim : q^{-1}(V) \in \mathcal{T}_X\}.$$

We now give an informal definition of a *cell complex*, for the purposes of this course.

Definition 1.5.8 (Disk) *We define the closed n -disk to be*

$$\mathbb{D}^n := \{x \in \mathbb{R}^n : d(0, x) \leq 1\} = \overline{B}(0, 1).$$

Per this definition, $\mathbb{D}^0 = \{\text{pt}\}$. Similarly, $\mathbb{D}^1 = [-1, 1]$. Now, in each of $n = 0, 1, 2, \dots$, take several copies of $\mathbb{D}^0, \mathbb{D}^1, \mathbb{D}^2, \dots$. In general, the boundary of \mathbb{D}^n is \mathbb{S}^{n-1} . We now form a process of “gluing” each of our copies to one another. In particular, we identify the boundary of a copy of

\mathbb{D}^n with a copy of \mathbb{D}^{n-1} . This is how we form the equivalence relation on the disjoint union of these discs. We call the quotient of the disjoint union, in this form, a *cell complex*.²⁷

27: Take 525 for a better definition.

Example 1.5.2 Going back to $[0, 1]$, if we take one copy of \mathbb{D}^0 and one copy of \mathbb{D}^1 , we glue the boundary of \mathbb{D}^1 ; i.e., $\{0, 1\}$, to the single point in \mathbb{D}^0 . Then, the rest of the equivalence classes are singletons.

Definition 1.5.9 (Saturated Set) Let X be a set and \sim an equivalence relation. A subset $U \subseteq X$ is called saturated with respect to \sim if for all $x, x' \in X$, then if $x \in U$ and $x' \sim x$, then $x' \in U$.

Using the definition of saturated set, we can reformulate our quotient topology by saying²⁸

$$\mathcal{T}_{X/\sim} = \{q(U) : U \text{ is saturated and } U \in \mathcal{T}_X\}.$$

28: Check this equivalence.

1.6 Homeomorphisms and Continuity

We now introduce the notion of a *homeomorphism*, which is an isomorphism in the category Top of topological spaces.²⁹

29: The morphisms in Top are precisely continuous maps.

Definition 1.6.1 (Homeomorphism) Given topological spaces (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) , a homeomorphism is a bijection $f : X \rightarrow Y$ such that³⁰

- (i) for all $U \in \mathcal{T}_X$, $f(U) \in \mathcal{T}_Y$.
- (ii) for any $V \in \mathcal{T}_Y$, $f^{-1}(V) \in \mathcal{T}_X$.

30: That is, $U \in \mathcal{T}_X$ if and only if $f(U) \in \mathcal{T}_Y$.

Remark 1.6.1 If there exists a homeomorphism $\varphi : X \rightarrow Y$ between spaces, then we say X is “homeomorphic” to Y , and write $X \simeq Y$.

How does one show that two topological spaces X and Y are *not* homeomorphic? Consider \mathbb{S}^1 and \mathbb{R} . We have that $\mathbb{S}^1 \not\simeq \mathbb{R}$, but how can we say that no such homeomorphism exists? Well, we need to build some topological *invariants* to distinguish our spaces.

Definition 1.6.2 (Projective Plane I) Start with \mathbb{S}^2 . Describe \sim on \mathbb{S}^2 by: for all $x \in \mathbb{S}^2$, $x \sim -x$. Then, the equivalence classes are $[x] = \{x, -x\}$, so the quotient space is

$$\mathbb{S}^2/\sim = \{[x] : x \in \mathbb{S}^2\}.$$

We denote $\mathbb{RP}^2 := \mathbb{S}^2/\sim$.

Definition 1.6.3 (Projective Plane II) Start with a copy of \mathbb{D}^2 and a copy of \mathbb{S}^1 . Well, $\partial\mathbb{D}^2 = \mathbb{S}^1$. Then, we can map $\partial\mathbb{D}^2 \rightarrow \mathbb{S}^1$, where half of $\partial\mathbb{D}^2$ gets sent to the complete circle \mathbb{S}^1 . This gives us our equivalence relation \sim' .³¹ Then, $\mathbb{RP}^2 := (\mathbb{D}^2 \sqcup \mathbb{S}^1)/\sim'$.

31: That is, antipodal points in $\partial\mathbb{D}^2$ are in the same equivalence class as a single point in \mathbb{S}^1 . Every point on the interior of \mathbb{D}^2 has singleton equivalence class.

32: This is very similar to our second definition. Constructing a homeomorphism between the first and second definition is a bit harder.

33: As an exercise, give five different definitions of continuous.

34: Any open set containing the given point is called a *neighborhood*.

35: This is with respect to the standard definition, via preimages.

36: The condition (ii), for a single $x \in X$, is called *continuity at the point x* .

Definition 1.6.4 (Projective Plane II') Start with a copy of \mathbb{D}^0 , \mathbb{D}^1 , and \mathbb{D}^2 . Identify the endpoints of \mathbb{D}^1 to the single point in \mathbb{D}^0 . Then, glue a pair of antipodal points in \mathbb{D}^2 to the same equivalence class. As before, half of \mathbb{D}^2 traverses the interval, and then the other half traverses the same interval in the same direction. This cell complex is $\mathbb{R}\mathbb{P}^2$.³²

Definition 1.6.5 (Continuous) Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces. A function $f : X \rightarrow Y$ is continuous if for all $V \in \mathcal{T}_Y$, the preimage $f^{-1}(V) \in \mathcal{T}_X$.³³

Remark 1.6.2 That is, a homeomorphism is a continuous bijection whose inverse is also continuous.

Lemma 1.6.1 Given $\mathcal{T}, \mathcal{T}'$ on X , the following are equivalent:

- (i) $\mathcal{T} \subseteq \mathcal{T}'$.
- (ii) for all $U \in \mathcal{T}$ and $x \in U$, there exists $V \in \mathcal{T}'$ such that $x \in V \subseteq U$.³⁴

Proof. For (i) \Rightarrow (ii), suppose $\mathcal{T}' \supseteq \mathcal{T}$. Let $V := U$. Now, for (ii) \Rightarrow (i), let $U \in \mathcal{T}$. Then, by (ii), for any $x \in U$, there exists $V_x \in \mathcal{T}'$ such that $x \in V_x \subseteq U$. Then, $U = \bigcup_{x \in U} V_x \in \mathcal{T}'$. \square

Lemma 1.6.2 Suppose $\mathcal{T}, \mathcal{T}'$ are topologies on X , and they are generated by bases $\mathcal{B}, \mathcal{B}'$, respectively. Then, the following are equivalent:

- (i) $\mathcal{T} \subseteq \mathcal{T}'$.
- (ii) for all $B \in \mathcal{B}$ and $x \in B$, there exists $B' \in \mathcal{B}'$ such that $x \in B' \subseteq B$.

Proof. The proof is left as an exercise. \square

We can state something similar with continuous functions.

Lemma 1.6.3 Let X, Y be topological spaces, and $f : X \rightarrow Y$ be some function. Then, the following are equivalent:

- (i) f is continuous.³⁵
- (ii) for all $x \in X$, and for any $V \in \mathcal{T}_Y$, if $f(x) \in V$, then there exists $U \in \mathcal{T}_X$ such that $x \in U$ and $f(U) \subseteq V$.³⁶

Sketch of Proof. Let us prove (ii) \Rightarrow (i). We want to show f is continuous, so take $V \in \mathcal{T}_Y$. Then, consider $f^{-1}(V) \subseteq X$. Then, for all $x \in f^{-1}(V)$, there exists $U_x \in \mathcal{T}_X$ such that $x \in U_x \subseteq f^{-1}(V)$, so $f^{-1}(V) = \bigcup_{x \in f^{-1}(V)} U_x$, so $f^{-1}(V) \in \mathcal{T}_X$. Thus, f is continuous. \square

Let us do one more of these.

Lemma 1.6.4 Let (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) be topological spaces, $\mathcal{B}_X, \mathcal{B}_Y$ be bases generating the respective topologies. Let $f : X \rightarrow Y$ be a function. Then, the

following are equivalent:

- (i) f is continuous.
- (ii) for all $x \in X$, for all $B' \in \mathcal{B}_Y$, if $f(x) \in B'$, then there exists $B \in \mathcal{B}_X$ such that $x \in B$ and $f(B) \subseteq B'$.³⁷

We could also state the definition of continuity in terms of closed sets: if $C \subseteq Y$ is closed, then $f^{-1}(C)$ is closed in X .

37: Once again, (ii) for a single $x \in X$, is precisely *continuity at the point* in terms of bases. Ranging over all x , we get an equivalent definition of continuity. When looking at a metric space, this is just ε, δ .

1.7 Product Topology

We now want to define a topology on the (cartesian) product of some topological spaces. If (X_1, \mathcal{T}_1) and (X_2, \mathcal{T}_2) are topological spaces, then define $X := X_1 \times X_2$. We can describe a set

$$\mathcal{B} := \{U_1 \times U_2 : U_1 \in \mathcal{T}_1, U_2 \in \mathcal{T}_2\}.$$

We want to know if we can squeeze something in the intersection of $U_1 \times U_2$ and $U'_1 \times U'_2$. Well, if $x \in (U_1 \times U_2) \cap (U'_1 \times U'_2)$, then

$$x \in (U_1 \cap U'_1) \times (U_2 \cap U'_2) \subseteq (U_1 \times U_2) \cap (U'_1 \times U'_2).$$

Thus, \mathcal{B} is a basis, so put the topology $\mathcal{T}(\mathcal{B})$ on X .³⁸

Let $\{(X_i, \mathcal{T}_i)\}_{i \in I}$ be a family (finite, countable, or uncountable) of topological spaces.

38: For a product of n spaces, we have a natural way to place a topology on the product. But, what happens if we have an infinite family of spaces?

Definition 1.7.1 (Arbitrary Cartesian Product) *We define the cartesian product*

$$X := \prod_{i \in I} X_i := \left\{ \underline{x} : I \rightarrow \bigcup_{i \in I} X_i : \text{for all } j \in I, \underline{x}(j) \in X_j \right\}.$$

Before we define a topology on this product, suppose we have a function

$$\begin{aligned} Y &\xrightarrow{f} \prod_{i \in I} X_i \\ y &\longmapsto \left(\underline{x} : I \rightarrow \bigcup_{i \in I} X_i \right), \end{aligned}$$

where Y is a topological space. We could define $f_j : y \mapsto f(y)(j) \in X_j$, where we picked some $j \in I$.³⁹ Each f leads to a family of functions $\{f_j : Y \rightarrow X_j\}_{j \in I}$.

39: This is *akin* to a restriction to the j th coordinate.

Definition 1.7.2 (Coordinate Function) *The function f_j , as above, is called the j th coordinate function.*

Conversely, each indexed family $\{f_j : Y \rightarrow X_j\}_{j \in I}$ yields a function $f : Y \rightarrow \prod X_i$, where $f(y)(j) := f_j(y)$.⁴⁰ In other words, f is entirely determined by its coordinate functions, and vice-versa. As such, we want

40: This new function f , is precisely the f we had originally.

a way to relate the continuity of f with the continuity of its coordinate functions by picking the needed topology on the product.

Consider the *canonical* projections

$$\begin{array}{ccc} \prod_{i \in I} X_i & \xrightarrow{\pi_j} & X_j \\ \underline{x} & \longmapsto & \underline{x}(j). \end{array}$$

Using this projection, we can define f_j as the composition given by the diagram

$$\begin{array}{ccc} Y & \xrightarrow{f} & \prod_{i \in I} X_i \\ & \searrow f_j := \pi_j \circ f & \downarrow \pi_j \\ & & X_j \end{array}$$

Figure 1.1: The existence of a unique f such that the diagram commutes is precisely the universal property of the product in **Set**. It turns out that via our construction of the product topology, we also get the universal property in **Top**.

Let $U_j \subseteq X_j$ and $U_j \in \mathcal{T}_j$. Then, the canonical preimage of U_j is

$$\pi_j^{-1}(U_j) = \left\{ \underline{x} : I \rightarrow \bigcup_{i \in I} X_i : \begin{array}{l} \text{for all } j \in I, \underline{x}(j) \in X_j \\ \text{and } \underline{x}(j) \in U_j \end{array} \right\}.$$

41: Why is this not a basis. Well consider $\pi_j^{-1}(U_j)$ and $\pi_{j'}^{-1}(U_{j'})$. Then, we should be able to squeeze in such a family which contains an \underline{x} in the intersection. This is not, generally possible.

Consider the family⁴¹

$$\mathcal{S} := \{ \pi_j^{-1}(U_j) : j \in I, U_j \in \mathcal{T}_j \}.$$

42: Take any $\underline{x} \in \prod_{i \in I} X_i$. Then, $\underline{x} \in \pi_j^{-1}(X_j)$, by definition.

Proposition 1.7.1 *The family \mathcal{S} , as above, is a subbasis.*⁴²

Definition 1.7.3 (Product Topology) *The product topology on $\prod X_i$ is $\mathcal{T}(\mathcal{B}(\mathcal{S}))$, the topology generated by the basis generated by the subbasis*

Remark 1.7.1 You will also hear the product topology referred to as the *Tychonoff* topology. It is the “most natural” one. One “proof” of this is via the *universal property*.

Remark 1.7.2 To assume that the arbitrary cartesian product of nonempty sets is nonempty is equivalent the axiom of choice.

It is worthwhile to mention that whereas $\mathcal{T}_{\text{prod}}$, as we have constructed it, picks finitely many indices in the basis. There is another topology on $\prod_i X_i$ known as the *box topology*.

Definition 1.7.4 (Box Topology) *The box topology on $\prod_i X_i$ is the topology*

$\mathcal{T}_{\text{box}} := \mathcal{T}(\mathcal{B}_{\text{box}})$ generated by the basis⁴³

$$\mathcal{B}_{\text{box}} := \left\{ \prod_{i \in I} U_i : \text{for all } i \in I, U_i \in \mathcal{T}_i \right\}.$$

43: The box topology is finer than the Tychonoff topology. Check this as an exercise.

1.8 Closure, Interior, and Boundary

Definition 1.8.1 (Closure of Subset) Let (X, \mathcal{T}_X) be a topological space and $A \subseteq X$. Then, the closure \bar{A} w.r.t. (X, \mathcal{T}_X) is defined as the intersection⁴⁴

$$\bar{A}^X := \bigcap_{A \subseteq C} C.$$

44: The $C \subseteq X$ are subsets which are closed with respect to \mathcal{T}_X and contain A .

Lemma 1.8.1 If $C \subseteq X$, then the following are equivalent:

- (i) C is closed in X .
- (ii) $\bar{C}^X = C$.

Proof. For (i) \Rightarrow (ii), we have that

$$\bar{C}^X = \bigcap_{C \subseteq C'} C' = C,$$

as $C \subseteq C$ is closed. Conversely, if $C = \bar{C}$, then

$$C^C = \left(\bigcap_{C \subseteq C'} C' \right)^C = \bigcup_{C \subseteq C'} (C')^C,$$

but the C' are closed, meaning $(C')^C \in \mathcal{T}_X$, so C^C is a union of open sets; i.e., its complement is open.⁴⁵ □

45: As such, we can characterize closedness via the closure.

Proposition 1.8.2 A point $x \in X$ does not belong to \bar{A}^X if and only if there exists an open neighborhood U of x such that $U \cap A = \emptyset$.

Proof. Begin with (\Leftarrow) . We have that U^C is closed, and $A \subseteq U^C$, so $\bar{A} \subseteq U^C \not\ni x$. Now, for (\Rightarrow) , suppose $x \notin \bar{A} = \bigcap_{A \subseteq C} C$. Pick any C like this⁴⁶ such that $x \in C^C$. □

46: We know the intersection is nonempty, as it does not contain x .

Remark 1.8.1 (Alternative Definition of Closure) Via the proposition, we have that $x \in \bar{A}$ if and only if for all open neighborhoods U of x , $U \cap A \neq \emptyset$. Thus,

$$\bar{A} := \{x \in X : \text{for all neighborhoods } U \text{ of } x, U \cap A \neq \emptyset\}.$$

Remark 1.8.2 The closure \bar{A} is always closed.⁴⁷

47: The lemma above gives the characterization, trivially.

48: We have that the LHS is an intersection of a *bigger* family than the RHS, so it is smaller.

Now, given $A_1 \subseteq A_2 \subseteq X$, what can we say about $\overline{A_1}$ and $\overline{A_2}$? Well, $\overline{A_1} \subseteq \overline{A_2}$. This is known as the *monotonicity of the closure*.⁴⁸

Lemma 1.8.3 Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$ a function. Then, the following are equivalent:

- (i) f is continuous.
- (ii) for any $A \subseteq X, f(\overline{A}) \subseteq \overline{f(A)}$.

49: That is, $f^{-1}(C) \subseteq X$ is closed, for any closed $C \subseteq Y$.

Proof. Using the closed set characterization of continuity, suppose f is continuous.⁴⁹ Take any set $A \subseteq X$. Let $x \in \overline{A}$. We want to show that $f(x) \in \overline{f(A)}$. Take any neighborhood V of $f(x)$. Since $f^{-1}(V)$ is open in X , and $x \in f^{-1}(V)$, we have that $A \cap f^{-1}(V) \neq \emptyset$, so $f(A) \cap f(f^{-1}(V)) \neq \emptyset$. As such, $f(A) \cap V \neq \emptyset$. Conversely, assume that for all $A \subseteq X, f(\overline{A}) \subseteq \overline{f(A)}$. Take any closed set $B \subseteq Y$. It will suffice to show that $f^{-1}(B)$ is closed in X ; i.e., that $f^{-1}(B) = \overline{f^{-1}(B)}$. Well, we already know that $f^{-1}(B) \subseteq \overline{f^{-1}(B)}$. By assumption,

$$f(\overline{f^{-1}(B)}) \subseteq \overline{f(f^{-1}(B))} \subseteq \overline{B} = B,$$

using the monotonicity of the closure and the closedness of B . Taking the preimage, we get

$$\overline{f^{-1}(B)} \subseteq f^{-1}(B),$$

as desired. □

50: Being Hausdorff gives us a notion of “separating” any two points. Morally, this means we would need a *fine enough* topology.

Definition 1.8.2 (Hausdorff) A topological space (X, \mathcal{T}) is Hausdorff if for all $x, y \in X$, there exists $U, V \in \mathcal{T}$ such that $x \in U$ and $y \in V$, and $U \cap V = \emptyset$.⁵⁰

Exercise 1.8.1 Let X, Y be Hausdorff. Show that $X \times Y$ equipped with $\mathcal{T}_{\text{prod}}$ is also Hausdorff. Conclude that $X \times Y$ equipped with \mathcal{T}_{box} is Hausdorff.

Definition 1.8.3 (Interior) Let (X, \mathcal{T}) and $A \subseteq X$. Then, we define the interior of A in X w.r.t. \mathcal{T} by

$$\text{int } A = \bigcup_{U \subseteq A} U,$$

where $U \in \mathcal{T}$.

Remark 1.8.3 It is immediate from the definition that $\text{int } A \subseteq A \subseteq \overline{A}$.

Definition 1.8.4 (Boundary) We define the boundary of $A \subseteq X$ w.r.t \mathcal{T} by

$$\partial A := \overline{A} \setminus \text{int } A.$$

Definition 1.8.5 (Limit Point) Given (X, \mathcal{T}) and $A \subseteq X$, a limit point of A in X is a point $x \in X$ such that for any neighborhood U of x in X ,⁵¹

$$(A \setminus \{x\}) \cap U \neq \emptyset.$$

51: The condition of being a limit point is stronger than being in the closure.

Remark 1.8.4 For any limit point $x \in X$ of A , $x \in \bar{A}$.

Proof. The proof is trivial. □

Example 1.8.1 Let $A := \{a_0, a_1\} \subseteq \mathbb{R}^2$. Then, a_0 is not a limit point, because there exists an open neighborhood U of a_0 such that $a_1 \notin U$, so

$$(\{a_0, a_1\} \setminus \{a_0\}) \cap U = \emptyset.$$

Yet, $a_0 \in A \subseteq \bar{A}$, so it is in the closure.

It turns out, this is *precisely* the general phenomenon. Define A' to be the set of all limit points of A in X w.r.t. \mathcal{T} .

Lemma 1.8.4 $\bar{A} = A \cup A'$.⁵²

52: We now have another equivalent definition of the closure, and in turn, we could restate our definition of continuity using the RHS.

Proof. We have that $A \cup A' \subseteq \bar{A}$, trivially. Conversely, let $x \in \bar{A}$.

- ▶ If $x \in A$, we are done.
- ▶ Suppose $x \notin A$. Then, $A \setminus \{x\} = A$, and since $x \in \bar{A}$, for any neighborhood $U \in \mathcal{T}$ of x , $A \cap U \neq \emptyset$, completing the proof.

□

1.9 Aside on Norms and Sequences of Functions

Recall that the *Euclidean* metric on \mathbb{R}^n is⁵³

$$d_2(x, y) := \sqrt{\sum_{i=1}^n (x_i - y_i)^2} = \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}}.$$

53: The “2” coming from the squaring and 1/2 power.

Similarly,

$$d_1(x, y) := \sum_{i=1}^n |x_i - y_i|.$$

In general, our p th metric is⁵⁴

$$d_p(x, y) := \sqrt[p]{\sum_{i=1}^n |x_i - y_i|^p}.$$

54: We leave checking that the d_p are metrics if $p \in [1, \infty)$.

55: The geometrical interpretation of the unit ball in \mathbb{R}^2 for each of the d metrics is useful. In some sense, d_∞ acts like a limiting case of the other d_p .

Remark 1.9.1 (Infinity Metric) We define⁵⁵

$$d_\infty := \max_{1 \leq i \leq n} \{|x_i - y_i|\}.$$

Definition 1.9.1 (Norm) A norm on a vector space $(\mathcal{V}, \mathbb{F})$, such as \mathbb{R}^n , is a function $\|\cdot\| : \mathcal{V} \rightarrow [0, \infty)$ such that

- (0) for all $x \in \mathcal{V}$, $\|x\| \geq 0$.
- (i) for all $x \in \mathcal{V}$, $\|x\| = 0$ if and only if $x = 0$.
- (ii) for all $x \in \mathcal{V}$ and $c \in \mathbb{F}$, $\|cx\| = |c| \cdot \|x\|$.
- (iii) for all $x, y \in \mathcal{V}$, $\|x + y\| \leq \|x\| + \|y\|$.

56: It is trivial to check that the properties of the norm imply the properties of the induced metric.

Now, given a norm $\|\cdot\| : \mathcal{V} \rightarrow [0, \infty)$ on $(\mathcal{V}, \mathbb{F})$, we can define an *induced distance function*⁵⁶

$$\begin{aligned} \mathcal{V} \times \mathcal{V} &\xrightarrow{d} [0, \infty) \\ (x, y) &\longmapsto \|x - y\|. \end{aligned}$$

Definition 1.9.2 (ℓ^2 -Norm) We define the ℓ^2 -norm on \mathbb{R}^n by

$$\begin{aligned} \mathbb{R}^n &\xrightarrow{\|\cdot\|_2} [0, \infty) \\ x &\longmapsto \sqrt{\sum_{i=1}^n x_i^2}. \end{aligned}$$

57: We are simply tracing our steps backward through the metrics.

Following this method, we can define the ℓ^p -norm on \mathbb{R}^n .⁵⁷

Definition 1.9.3 (ℓ^p -Norm) Generally, the ℓ^p -norm on \mathbb{R}^n is defined by

$$\begin{aligned} \mathbb{R}^n &\xrightarrow{\|\cdot\|_p} [0, \infty) \\ x &\longmapsto \sqrt[p]{\sum_{i=1}^n |x_i|^p}. \end{aligned}$$

Definition 1.9.4 (Real Sequence Space) The real sequence space is

$$\mathbb{R}^\omega = \mathbb{R}^{\mathbb{Z}_+} := \{f : \mathbb{Z}_+ \rightarrow \mathbb{R}\}.$$

58: On this set, $\|\cdot\|_2$ is a well-defined norm.

Definition 1.9.5 ($\ell^2(\mathbb{Z}_+, \mathbb{R})$ Sequence Space) Define⁵⁸

$$\ell^2(\mathbb{Z}_+, \mathbb{R}) := \left\{ f : \mathbb{Z}_+ \rightarrow \mathbb{R} : \sum_{i \in \mathbb{Z}_+} |f(i)|^2 < \infty \right\} \subseteq \mathbb{R}^\omega.$$

Definition 1.9.6 ($\ell^p(\mathbb{Z}_+, \mathbb{R})$ Sequence Space) *Following suit, define*

$$\ell^p(\mathbb{Z}_+, \mathbb{R}) := \left\{ f : \mathbb{Z}_+ \rightarrow \mathbb{R} : \sum_{i \in \mathbb{Z}_+} |f(i)|^p < \infty \right\}.$$

Here, the ℓ^p norm is precisely

$$\|f\|_p := \sqrt[p]{\sum_{i \in \mathbb{Z}_+} |f(i)|^p}$$

As is standard in the literature, we will denote $\ell^p := \ell^p(\mathbb{Z}_+, \mathbb{R})$. Then, we have an inclusion $\iota : \ell^1 \hookrightarrow \ell^2$.⁵⁹

Note that we can also define the ℓ^∞ norm via

$$\|f\|_\infty := \sup_{i \in \mathbb{Z}_+} \{|f(i)|\} \in [0, \infty] \subset \mathbb{R} \cup \{\infty\}.$$

In this sense,

$$\ell^\infty(\mathbb{Z}_+, \mathbb{R}) := \{f : \mathbb{Z}_+ \rightarrow \mathbb{R} : \|f\|_\infty < \infty\}.$$

Of course, $|\cdot|$ is a norm on \mathbb{R} , so $\|\cdot\|_\infty$ is a norm on ℓ^∞ .⁶⁰

Definition 1.9.7 (Lipchitz Equivalent Metrics) *Two metrics d, d' are Lipchitz equivalent on X if there exist $c_1, c_2 > 0$ such that*

$$d \leq c_1 d' \text{ and } d' \leq c_2 d.$$

Definition 1.9.8 (Lipchitz Function) *A function $f : (X, d_X) \rightarrow (Y, d_Y)$ is called Lipchitz if there exists a constant $L > 0$ such that for all $x, x' \in X$,*

$$d_Y(f(x), f(x')) \leq L d_X(x, x').$$

Definition 1.9.9 (Sequence Convergence) *A sequence $\{x_i\} \subseteq X$ on a topological space (X, \mathcal{T}) converges to $x \in X$ if for all neighborhoods U of x , there exists an $N \in \mathbb{Z}_+$ such that for all $n \geq N$, $x_n \in U$.*

Definition 1.9.10 (Sequentially Closed) *A subset $A \subseteq X$ is sequentially closed if for any sequence $\{x_i\} \subseteq A$ which converges to $x \in X$, then $x \in A$.⁶¹*

Lemma 1.9.1 *Let X be a topological space and $A \subseteq X$, then $\overline{A}^{\text{seq}} \subseteq \overline{A}$. That is to say, for any sequence $\{x_n\}$ in A with $x_n \rightarrow x \in X$, then $x \in \overline{A}$.*

Proof. Take any sequence $\{x_n\}$ in A such that $x_n \rightarrow x \in X$. Then, $x \in \overline{A}^{\text{seq}}$. Take any neighborhood U of x in X . Since $x_n \rightarrow x$, there exists $N \in \mathbb{Z}_+$ such that for all $n \geq N$, $x_n \in U$, but $x_n \in A$, so $U \cap A \neq \emptyset$. \square

What about the converse? In general, it is not true. However, we can make our space *nice enough* for it to hold.

59: The opposite inclusion certainly fails. Consider the harmonic sequence

$$f(i) = \frac{1}{i}.$$

We have that $f \notin \ell^1$, but $f \in \ell^2$.

60: Check this as an exercise.

61: Prove that our standard definition of being closed implies sequentially closed. However, the converse does not hold, generally.

Definition 1.9.11 (Metriizable) *A topological space (X, \mathcal{T}) is metriizable if there exists a metric d on X such that $\mathcal{T} = \mathcal{T}_d$.*

Lemma 1.9.2 *Let X be a metriizable topological space. Then, $\overline{A} \subseteq \overline{A}^{\text{seq}}$.*

Proof. Let $x \in \overline{A}$. Consider the ball $B(x, 1)$. Well, $B(x, 1) \cap A \neq \emptyset$. That is, there exists x_1 in the intersection. Similarly, consider the ball $B(x, 1/2) \cap A \neq \emptyset$, so we get an x_2 . In general, we can define a sequence $\{x_n\}$ in A via $B(x, 1/n)$.⁶² To show that $x_n \rightarrow x$, take any U of x . Then, there exists a ball B centered at x such that $B \subseteq U$. Let r be the radius of B . Let $N = 1/r + 1$. Then every x_n for $n \geq N$ is contained in B , so they are contained in U . \square

62: We use the axiom of choice here.

Definition 1.9.12 (Basis at Point) *A basis at $x \in X$ is a family of open neighborhoods $\{U_i\}_{i \in \Lambda}$ of x such that for any neighborhood U of x , there exists an $i \in \Lambda$ such that $x \in U_i \subseteq U$.*

Definition 1.9.13 (First-Countable Space) *A topological space (X, \mathcal{T}) is called first-countable if for any $x \in X$, there exists a countable basis at x .⁶³*

63: All metriizable spaces are first-countable, as we can take a basis of open balls of radius $1/n$.

If we have a countable basis U_1, U_2, \dots , then we can define $U'_1 := U_1$, $U'_2 := U_1 \cap U_2$, $U'_3 := U_1 \cap U_2 \cap U_3$, and so forth.⁶⁴

64: Note that a second-countable space is one with a countable generating basis.

Proposition 1.9.3 *Let X be a first-countable topological space. Then, $\overline{A} \subseteq \overline{A}^{\text{seq}}$.*

Theorem 1.9.4 (Sequentially Continuous) *A function $f : X \rightarrow Y$ is sequentially continuous if for any sequence $\{x_n\}$ in X and any x in X , if $x_n \rightarrow x$, then $f(x_n) \rightarrow f(x)$.⁶⁵*

65: That is, f maps convergent sequences (and in particular, the limit) to convergent sequences (image of the limit).

We can now ask, what is the relationship between sequential continuity and our standard continuity.

Lemma 1.9.5 *Let $f : X \rightarrow Y$ be a function between topological spaces. If X is continuous, then it is sequentially continuous. If we additionally have that X is first-countable (notably, metriizable), then the converse holds.*

Proof. Start with (\Rightarrow) . Suppose $f : X \rightarrow Y$ is continuous. Take any sequence $\{x_n\} \subseteq X$ such that $x_n \rightarrow x \in X$. Take any open neighborhood V of $f(x)$. Then, $f^{-1}(V) \in \mathcal{T}_X$. Then, there exists an $N \in \mathbb{Z}_+$ such that $x_n \in f^{-1}(V)$ for all $n \geq N$, so $f(x_n) \in V$, meaning $f(x_n) \rightarrow f(x)$. Conversely, assume X is first-countable. Then, for any $A \subseteq X$, $\overline{A} = \overline{A}^{\text{seq}}$. Take a closed $B \subseteq Y$. We have that $f^{-1}(\overline{B}) = \overline{f^{-1}(B)}^{\text{seq}}$. Let $x \in \overline{f^{-1}(B)}^{\text{seq}}$. Then, there exists a sequence $\{x_n\} \subseteq f^{-1}(B)$ such that $x_n \rightarrow x$. Then, $f(x_n) \rightarrow f(x) \in \overline{B} = B$. Thus, $x \in f^{-1}(B)$, so $\overline{f^{-1}(B)} = f^{-1}(B)$, meaning $f^{-1}(B)$ is closed. \square

Now, note that

$$\text{second-countable} \Rightarrow \text{first-countable} \Leftarrow \text{metrizable}.$$

Take countably many copies of $[0, 1]$, take their disjoint union, and form the cell complex gluing all the 0s together. We will see this space later on, but it is worth noting that this graph is *not first-countable* with the quotient topology.⁶⁶ We will prove the result later on.

66: Obviously, if you take the topology induced by the plane, it is first-countable.

Remark 1.9.2 In general, given a cell complex which is not *locally finite*, we can say that it is not first-countable or metrizable via the quotient topology.

Consider the set of all possible functions $\mathfrak{F}(\mathbb{R}, \mathbb{R}) = \{f : \mathbb{R} \rightarrow \mathbb{R}\}$. How can we put a topology on this set? Let us generalize a bit. Let $\{f_n\}$ be a sequence of functions in $\mathfrak{F}(X, Y)$, where X, Y are fixed topological spaces, Let $f : X \rightarrow Y$ be another function.

Definition 1.9.14 (Pointwise Convergence) We say $f_n \rightarrow f$ pointwise if for all $x \in X$, $f_n(x) \rightarrow f(x)$.⁶⁷

67: We are using the well-defined notion of convergence in Y .

Suppose additionally that (Y, d_Y) is a metric space. Then, $f_n \rightarrow f$ pointwise is equivalent to: for all $x \in X$, for any $\varepsilon > 0$, there exists $N \in \mathbb{Z}_+$ so that for $n \geq N$, $f_n(x) \in B(f(x), \varepsilon)$. What if we make this *uniform*? For all $\varepsilon > 0$, there exists $N \in \mathbb{Z}_+$ such that for all $x \in X$ and $n \geq N$, $f_n(x) \in B(f(x), \varepsilon)$. The uniform converge in a metric space codomain is precisely taking

$$\sup_x \{d(f_n(x), f(x))\} \leq \varepsilon.$$

Definition 1.9.15 (Supremum Metric) Let $f, g : X \rightarrow Y$. Then,⁶⁸

$$d_{\text{sup}}(f, g) := \sup_x \{d_Y(g(x), f(x))\}.$$

68: Check that this is a distance function on the set of all bounded functions.

Definition 1.9.16 (Uniform Convergence) Let $f, f_n : X \rightarrow Y$, where X is a set and Y is a metric space. We say $f_n \rightarrow f$ uniformly if for all $\varepsilon > 0$, there exists an $N \in \mathbb{Z}_+$ such that for all $n \geq N$,

$$d_{\text{sup}}(f_n, f) \leq \varepsilon.$$

Theorem 1.9.6 (Uniform Limit Theorem) Let X be a topological space and Y a metric space, and $f_n : X \rightarrow Y$ is a sequence of continuous functions converging uniformly to a function $f : X \rightarrow Y$. Then, f is continuous.

Sketch of Proof. Let V be an open set $V \subseteq Y$. We wish to show that $f^{-1}(V)$ is open. Take any point $x \in f^{-1}(V)$. Consider $f(x)$. Pick a small enough $\varepsilon > 0$ so that $B(f(x_0), \varepsilon) \subseteq V$. Choose N large enough so that for any $x \in X$, the distance $d(f_n(x), f(x)) < \varepsilon/3$. Then, choose a neighborhood U of x_0 in X such that $f_N(U) \subseteq B(f_n(x_0), \varepsilon/3)$.⁶⁹ \square

69: We can do this step and the previous one via uniform convergence and continuity, respectively.

Connectedness and Compactness

2

We now consider some standard point-set topological invariants and characteristics. You likely have seen these in real analysis.

2.1 (Path-)Connectedness

Before we discuss connectedness, what would it mean for a space to *not* be connected? Well, $[0, 1] \cup [3, 4]$ is certainly not connected, intuitively.

Definition 2.1.1 (Separation) *A separation of a topological space X is a pair of open, nonempty subsets (U, V) in X such that $U \cap V = \emptyset$ and $X = U \cup V$.¹*

Definition 2.1.2 (Connected) *A topological space X is connected if it does not admit a separation (U, V) on X .²*

Definition 2.1.3 (Path-Connected) *A space X is path-connected if for any $x, y \in X$, there exists a continuous function*

$$\gamma : [0, 1] \rightarrow X$$

such that $\gamma(0) = x$ and $\gamma(1) = y$.³

Lemma 2.1.1 *Any path-connected space is connected.*

Proof. Suppose X is not connected. Then, there exists a separation (U, V) of X such that $X = U \cup V$ and $U \cap V = \emptyset$.⁴ Pick $x \in U$ and $y \in V$. Then, there exists a path γ from x to y . Then, $(\gamma^{-1}(U), \gamma^{-1}(V))$ is a separation of $[0, 1]$, a contradiction. \square

Define an equivalence relation \sim_p on a topological space X by $x \sim_p y$ if and only if there exists a path $\gamma : [0, 1] \rightarrow X$ so that $\gamma(0) = x$ and $\gamma(1) = y$.

Lemma 2.1.2 *The relation \sim_p is an equivalence.*⁵

Definition 2.1.4 (Path Component) *A path component is an equivalence class of \sim_p in X .*

Define another equivalence relation \sim on X by $x \sim y$ if and only if there exists a connected subset $A \subseteq X$ such that $x, y \in A$. Take the subspace topology on A .

2.1 (Path-)Connectedness	21
2.2 Compactness	24
2.3 Local and Sequential Compactness	26
2.4 Lebesgue's Number Lemma	28
2.5 Compactifications and Zorn's Lemma	30

1: The intuitive example admits a separation, by construction.

2: Read the Munkres proof of connectedness of intervals. It is not too difficult.

3: We call any such map γ a *path* in X .

4: Recall that $U, V \neq \emptyset$.

5: Use the constant map for reflexivity. For symmetry, pre-compose γ with $f(t) = 1 - t$ to get $\bar{\gamma}$ in the form we desire. For transitivity, concatenate the paths (though, proving continuity is a bit technical).

6: Reflexivity comes from $A = \{x\}$. Symmetry is trivial. Transitivity is done via the union and the following lemma.

7: Recall that U, V are open in A via the subspace topology.

8: The pre-images are nonempty, as $\gamma(0) = x$ and $\gamma(1) = y$.

9: Use the same argument as above.

10: If X is path-connected via γ and $f : X \rightarrow Y$ is continuous, then $f \circ \gamma$ is a path for Y .

Lemma 2.1.3 *The relation \sim is an equivalence.*⁶

Lemma 2.1.4 *Let X be a topological space and $\{A_i\}_{i \in I}$ a collection of connected subsets $A_i \subseteq X$. Suppose there exists a point $x_0 \in X$ such that $x_0 \in \bigcap A_i$. Then, $\bigcup A_i$ is connected.*

Proof. Suppose $A := \bigcup A_i$ is not connected. Then, there exists a separation (U, V) of A .⁷ Well, $x_0 \in U \cup V$, so without loss of generality we can assume $x_0 \in U$, as U, V are disjoint. For each index $i \in I$, we know $x_0 \in A_i \subseteq U \cup V$. Yet, A_i is connected, so $A_i \subseteq U$ or $A_i \subseteq V$, and since $x_0 \in U$, $A_i \subseteq U$. This works for every $i \in I$, so $V = \emptyset$, a contradiction. \square

Definition 2.1.5 (Connected Component) *A connected component is an equivalence class of \sim in X .*

Definition 2.1.6 (Locally Path-Connected) *A space X is locally path-connected if for all $x \in X$ and for all open neighborhoods U of x , there exists another neighborhood U' of x such that $x \in U' \subseteq U$ and U' is path-connected.*

Definition 2.1.7 (Locally Connected) *A space X is locally connected if for all $x \in X$ and for all open neighborhoods U of x , there exists another neighborhood U' of x such that $x \in U' \subseteq U$ and U' is connected.*

Remark 2.1.1 Use the subspace topology for U , for the local definitions.

Lemma 2.1.5 *If $x \sim_p y$ then $x \sim y$.*

Proof. Let γ be the path for $x \sim_p y$. Consider $\gamma([0, 1]) \subseteq X$. Suppose $\gamma([0, 1])$ admits a separation (U, V) . Then, pre-images give us a contradiction, so $\gamma([0, 1])$ is connected, meaning $x \sim y$.⁸ \square

Corollary 2.1.6 *The continuous image of any connected space is connected.*⁹

Proposition 2.1.7 *The continuous image of any path-connected space is path-connected.*¹⁰

Theorem 2.1.8 *If X is any topological space, $A \subseteq X$ is connected, and $f : X \rightarrow Y$ is continuous, then $f(A)$ is connected.*

From our discussion above, it is clear that for all $x \in X$, $[x]_{\sim_p} \subseteq [x]_{\sim}$. That is, the partition of X generated by \sim_p is finer than that by \sim .

Lemma 2.1.9 *A space X is connected if and only if it has exactly one connected component.*

Proof. Suppose X is connected. Let $x \in X$. For each $x' \in X$, $x, x' \in X$ and X is connected. Since x' is arbitrary, $x \sim x'$, so $X \subseteq [x]_{\sim}$. Thus, $X = [x]_{\sim}$, since there is nothing larger. Conversely, assume there exists exactly one connected component: $\bigcup [x]_{\sim}$ is connected, but this is all of X . \square

Theorem 2.1.10 *If a topological space X is connected and locally path-connected, then it is path-connected.*

Proof. We want to show that X has precisely *one* path component. Since X is connected, there exists exactly one connected component: X . The path components give a partition of X . Let $x \in X$ and consider the path component $[x]_{\sim_p}$. Since the space is locally path-connected, there exists a path-connected neighborhood U'_x of x in X , so $U'_x \subseteq [x]_{\sim_p}$. Do this for all $x' \in [x]_{\sim_p}$. Thus, $\bigcup U'_{x'} = [x]_{\sim_p}$, so each path component of X is open. If there exists at least 2 path components, then we have a separation of X , which is a contradiction.¹¹ \square

11: We use that we can union the path components to form a nonempty, open, disjoint partition.

Remark 2.1.2 On the way to proving the theorem, we showed that if a space is locally path-connected, then the path components are open. The same implication holds for connectedness.

Lemma 2.1.11 *For any connected subset $A \subseteq X$, if $A \subseteq B \subseteq \bar{A}$, then B is connected.*¹²

12: In particular, \bar{A} is closed

Now, let $C \subseteq X$ be a connected component. Then, C is indeed connected, so \bar{C} is connected. Thus, $\bar{C} = C$, so C is closed.

Theorem 2.1.12 (Intermediate Value Theorem) *If X is a connected topological space and $f : X \rightarrow \mathbb{R}$ is continuous in the standard topology, then for any $x_0, x_1 \in X$ and $c \in \mathbb{R}$ between $f(x_0)$ and $f(x_1)$, inclusive, then there exists $x \in X$ so that $f(x) = c$.*

Proof. Suppose not. Then, $c \notin f(X)$. Then, the image $f(X) \subseteq (-\infty, c) \cup (c, \infty)$. Both $f(X) \cap (-\infty, c)$ and $f(X) \cap (c, \infty)$ are open by the subspace topology, so the intersections form a separation of $f(X)$. Take the pullback. Then, $f^{-1}(-\infty, c)$ and $f^{-1}(c, \infty)$ form a separation of X .¹³ \square

13: This is a contradiction. The standard application is for intervals in \mathbb{R} .

Remark 2.1.3 (Topologist's Sine Curve) Union the graph of $\sin(1/x)$ in \mathbb{R}^2 to the interval on the y -axis. This is connected, but you get a nice contradiction to the continuity of any path from the interval to $\sin(1/x)$.

Remark 2.1.4 Note that if a space is locally path-connected, then the path components are open, so we could form a separation of a surrounding connected component. Hence, path components and connected components coincide exactly when the whole space is locally path-connected.¹⁴

14: This is another way to find the $x \sim y \Rightarrow x \sim_p y$ implication.

Theorem 2.1.13 $\mathbb{R} \not\cong \mathbb{S}^1$.

Proof. Let $\varphi : \mathbb{S}^1 \xrightarrow{\sim} \mathbb{R}$ be a homeomorphism. Let $a \in \mathbb{S}^1$, and suppose $f(a) = b \in \mathbb{R}$. Consider $\mathbb{S}^1 \setminus \{a\}$ and $\mathbb{R} \setminus \{b\}$. Then, $\mathbb{S}^1 \setminus \{a\}$ is path-connected, but there is an obvious separation of $\mathbb{R} \setminus \{b\}$.¹⁵ \square

15: Do this rigorously with a restriction.

2.2 Compactness

We now want to establish a sense of “smallness” for arbitrary topological spaces, lacking a notion of boundedness.

Definition 2.2.1 (Cover) *A cover of a set X is a family $\{U_i \subseteq X\}_{i \in I}$ so that the union $\bigcup U_i = X$.*

If (X, \mathcal{T}) is a topological space, then an *open cover* of X is a cover $\{U_i\}_{i \in I}$ such that $U_i \in \mathcal{T}$ for all $i \in I$.

Definition 2.2.2 (Compact) *A topological space X is called compact if for any open cover $\{U_i\}_{i \in I}$ of X , there exists a finite subcover of X .*

Example 2.2.1 Consider $(0, 1) \simeq \mathbb{R}$. Consider the family $\{(i, i + 2)\}_{i \in \mathbb{Z}}$. Then, $\bigcup (i, i + 2) = \mathbb{R}$. Any finite subfamily would have a smallest and largest i . Then, $(i_{\min}, i_{\max} + 2) \neq \mathbb{R}$, so \mathbb{R} is not compact, meaning $(0, 1)$ is not either.

Example 2.2.2 (Empty Set) \emptyset is compact, vacuously.

Example 2.2.3 (Singleton) Singletons $\{x\} \subset \mathbb{R}$ are compact.

Example 2.2.4 (Closed Interval) Consider $[0, 1] \subset \mathbb{R}$. We claim it is compact.

We might call the original definition of compactness the *intrinsic* definition. What would a *relative* or *extrinsic* definition be?

Definition 2.2.3 (Subset Compactness) *Let (X, \mathcal{T}_X) be a topological space and $C \subseteq X$. We call C compact if for any family $\{U_i \in \mathcal{T}_X\}_{i \in I}$ so that $C \subseteq \bigcup_I U_i$, then there is a finite subfamily $\{U_i \in \mathcal{T}_X\}_{i \in I'}$ so that $I' \subseteq I$ with $|I'| < \infty$ and $C \subseteq \bigcup_{I'} U_i$.*

16: Chasing definitions makes this an easy proof.

Theorem 2.2.1 *If $C \subseteq X$, the two definitions of compactness are equivalent.*¹⁶

Lemma 2.2.2 *If (X, \mathcal{T}_X) is a compact topological space, then any closed set $C \subseteq X$ is compact.*

Proof. Pick a “generalized” open cover $\mathcal{F} := \{U_i \in \mathcal{T}_X\}_{i \in I}$ so that $C \subseteq \bigcup_I U_i$. Now, $X \setminus C$ is open, so $\mathcal{F}' := \{U_i\}_{i \in I} \cup (X \setminus C)$ is an open cover of X . Since X is compact, there exists a finite subfamily $\{U_{i_1}, \dots, U_{i_n}\}$ of \mathcal{F}' that covers X . Thus, C is compact.¹⁷ \square

17: We use the second definition.

Lemma 2.2.3 *Let (X, \mathcal{T}_X) be Hausdorff and $C \subseteq X$ a compact subspace. Then, C is closed in X .*

Proof. We want to show $X \setminus C$ is open. Fix a point $y \in X \setminus C$. Since X is Hausdorff, for any $x \in C$, we can find neighborhoods U_x of x and V_x of y such that $U_x \cap V_x = \emptyset$. Do this for each choice of $x \in C$. Consider the family $\{U_x\}_{x \in C}$. Well, $C \subseteq \bigcup_C U_x$.¹⁸ Since C is compact, there exists a finite subcover $\{U_{x_1}, \dots, U_{x_n}\}$ of C . Consider the corresponding $\{V_{x_1}, \dots, V_{x_n}\}$. Then,

18: That is, $\{U_x\}$ is a generalized open cover of C .

$$\underbrace{\left(\bigcap_{i=1}^n V_{x_i}\right)}_V \cap \left(\bigcup_{i=1}^n U_{x_i}\right) = \emptyset,$$

so $C \cap V = \emptyset$ and $y \in V$. This can be done for every choice of $y \in X \setminus C$. Hence, $X \setminus C$ is open, meaning C is closed.¹⁹ \square

19: This is a nifty result. No matter how you homeomorphically embed a compact space in a Hausdorff space, it will always be closed, despite the fact that closedness is relative to ambient topology.

Lemma 2.2.4 *Let $f : X \rightarrow Y$ be a continuous function between topological spaces and $C \subseteq X$ be a compact subspace. Then, $f(C) \subseteq Y$ is compact.*

Proof. Let $\{U_i \in \mathcal{T}_Y\}_{i \in I}$ be an open cover of $f(C)$. Clearly $f^{-1}(U_i)$ is open for all $i \in I$. Since the U_i cover $f(C)$, the $f^{-1}(U_i)$ cover C . There is a finite subset $I' \subseteq I$ such that $\{f^{-1}(U_i)\}_{i \in I'}$ is a cover of C . Hence, $\{U_i\}_{i \in I'}$ is an open cover of $f(C)$. \square

For any product $X \times Y$, we can pick any $x \in X$. Then, we can consider the slice

$$Y_x := \{(x, y) : y \in Y\} = \{x\} \times Y.$$

Define a map²⁰

$$\begin{aligned} Y &\xrightarrow{\varphi_x} Y_x \\ y &\longmapsto (x, y). \end{aligned}$$

20: We claim that φ_x is a bijection.

In fact, the function φ_x is a homeomorphism with the subspace topology on $Y_x \subseteq X \times Y$, which can be seen using open rectangles and projections.

Lemma 2.2.5 *If X and Y are compact topological spaces, then $X \times Y$ is compact with respect to the product topology.*

Proof. Fix $x \in X$. Consider the corresponding slice Y_x . Since Y is compact, and $Y \simeq Y_x$, we know Y_x is compact with respect to the subspace topology from the product topology. Let $\mathcal{A} := \{A_i\}_{i \in I}$ be an open cover of $X \times Y$.²¹ For any $y \in Y$, there exists an $i \in I$ such that $(x, y) \in A_i$. For each y , pick an open $U_{x,y} \subseteq X$ and open $V_{x,y} \subseteq Y$ such that $(x, y) \in U_{x,y} \times V_{x,y} \subseteq X \times Y$.

21: Our goal is to extract a finite cover.

The family $\{U_{x,y} \times V_{x,y}\}_{y \in Y}$ is an open cover of $\{x\} \times Y = Y_x$. Then, there exists a finite subcover $\{U_{x,y_1} \times V_{x,y_1}, \dots, U_{x,y_n} \times V_{x,y_n}\}$. Now,

$$\{x\} \times Y \subseteq \underbrace{\left(\bigcap_{i=1}^n U_{x,y_i} \right)}_{W_x \in \mathcal{O}_X} \times Y.$$

The set $\{W_x : x \in X\}$ is an open cover of X . Since X is compact, we can take finitely many W_{x_k} for $k \in [m]$. Then, $\{W_{x_k} \times Y\}_{k \in [m]}$ is a finite open cover of $X \times Y$. Tracing back through our steps, each tube is contained in one of the open rectangles, and each of the open rectangles is contained in one of the $A_i \in \mathcal{A}$. Since we have made finitely many choices of such A_i to cover $X \times Y$, we are done. \square

22: This works for all finite products, as you might expect.

Remark 2.2.1 There is a natural isomorphism in Set between $X \times Y \times Z \simeq (X \times Y) \times Z$, and taking the corresponding product topologies, we get that $X \times Y \times Z$ is compact.²²

Corollary 2.2.6 All finite products of compact sets are compact.

23: We omit the proof for now. We will come back to this, though. It really is a neat proof.

Theorem 2.2.7 (Tychonoff Theorem) If $\{X_i\}_{i \in I}$ is a family of connected topological spaces, then $\prod X_i$ is compact.²³

2.3 Local and Sequential Compactness

Theorem 2.3.1 (Extreme Value Theorem) Given a continuous function $f : X \rightarrow \mathbb{R}$, where X is a compact topological space, there exist $x_{\min}, x_{\max} \in X$ such that for all $x \in X$,

$$f(x_{\min}) \leq f(x) \leq f(x_{\max}).$$

Proof. Suppose that for all $x' \in X$, there exists $x \in X$ so that $f(x') < f(x)$. We have $f(X) \subseteq \mathbb{R}$. Now, consider the set

$$\{(-\infty, a) : a \in f(X)\}.$$

Then, $f(X) \subseteq \bigcup_{a \in f(X)} (-\infty, a)$. In particular, this set is an open cover of $f(X)$. Now, X is compact and f is continuous, so $f(X)$ is compact, which means there exists a finite subcover $\{(-\infty, a_i) : i \in [n]\}$ of $f(X)$. Thus, $f(X) \subseteq \bigcup (-\infty, a_i)$. Pick $a := \max\{a_1, \dots, a_n\} \in f(X)$. Then, $f(X) \subseteq (-\infty, a)$, and $a \notin (-\infty, a)$, so $a \notin f(X)$. This is a contradiction.²⁴ \square

24: We assume all the a are in $f(X)$.

Remark 2.3.1 The same process works for x_{\min} .

We will now define local compactness, which does *not* follow our usual pattern for locality.

Definition 2.3.1 (Locally Compact) *A space X is called locally compact if for any $x \in X$ there exists a neighborhood U of x which is contained in a compact subset $C \subseteq X$.²⁵*

25:
$$x \in U \subseteq C \subseteq X$$

Proposition 2.3.2 *Compactness trivially implies local compactness. The converse, however, is not true.*

Example 2.3.1 We know \mathbb{R} is not compact. Yet, \mathbb{R} is locally compact.²⁶

26: Take a closed interval C and a smaller open interval U .

Definition 2.3.2 (Subsequence) *A subsequence of $\{x_n\}$ given by $f : \mathbb{Z}_+ \rightarrow X$ is any composition*

$$\begin{array}{ccc} \mathbb{Z}_+ & \hookrightarrow & \mathbb{Z}_+ \xrightarrow{x} X \\ k & \longmapsto & n_k \end{array}$$

such that $n_k < n_{k'}$ if $k < k'$.

Definition 2.3.3 (Sequentially Compact) *A topological space X is sequentially compact if for any sequence $\{x_n\} \subseteq X$, there exists a subsequence $\{x_{n_k}\} \subseteq X$ such that $x_{n_k} \rightarrow L$ for some $L \in X$.*

Theorem 2.3.3 *Let X be metrizable (or first-countable). Then, the following are equivalent:*

- (i) X is compact.
- (ii) X is sequentially compact.

Proof. We will just show (i) \Rightarrow (ii).²⁷ Take any sequence $\{x_n\} \subseteq X$. Define $A := \{x_n : n \in \mathbb{Z}_+\} \subseteq X$. We defined that $y \in A'$ if and only if for any neighborhood U of y , $(A \setminus \{y\}) \cap U \neq \emptyset$. Start with the case $A' = \emptyset$. Then, $\overline{A} = A \cup A' = A$, so A is closed. Then, $X \setminus A$ is open. For any $a \in A$, the negation says there exists a neighborhood U_a of a such that $A \setminus a \cap U_a = \emptyset$, so $A \cap U_a = \{a\}$. Consider the family $\{U_a\}_{a \in A} \cup (X \setminus A)$. This is an open cover of X . Compactness tells us that there exists a finite subcover $\{U_{a_i}\}_{i=1}^m \cup \{X \setminus A\}$. That is, A is finite! From here you deduce that you can extract a convergent constant subsequence. Case two is when $A' \neq \emptyset$. Take $x \in A'$. Take an open ball U around x . We know $(A \setminus \{x\}) \cap U \neq \emptyset$. Successively, we take $B(x, 1), B(x, r_1)$, where $r_1 < 1$. Continue, taking smaller and smaller balls. Each time, we pick $y_i \in A$ with a larger index. We can check that the system of radii converges to zero, meaning we have a local basis at x . Hence, $y_i \rightarrow x$ in the sense of topology.²⁸ \square

27: Read Munkres for the proof of the converse.

28: Take any neighborhood of x , then at some point, one of the balls in our local basis will fit in.

Definition 2.3.4 (Bounded) *A metric space (X, d) is bounded if the diameter*

$$\text{Diam}(X) := \sup\{d(x_1, x_2) : x_1, x_2 \in X\} < \infty$$

29: The bound comes from the triangle inequality.

Remark 2.3.2 In \mathbb{R}^n , boundedness of C amounts to picking a ball $B(x, r)$ so that $C \subset B(x, r)$, where the diameter $\text{Diam}(X)$ is bounded by $2r$.²⁹

Theorem 2.3.4 (Heine-Borel) *For any subset $C \subseteq \mathbb{R}^n$, C is compact if and only if C is closed and bounded.*

Proof. Suppose C is compact. Since \mathbb{R}^n is Hausdorff, C is closed. Consider the set $\{B(x, 1) : x \in C\}$. This forms an open cover of C . We have a finite subcover $\{B(x_i, 1) : 1 \leq i \leq n\}$ of C . Let $\mathcal{D} := \max\{d(x_1, x_i) : 1 \leq i \leq n\}$. Then, $C \subseteq B(x, \mathcal{D} + 2)$, so C is bounded. Conversely, suppose $C \subseteq \mathbb{R}^n$ is closed and bounded. There exists a cube $Q = [a, b]^n$ so that $C \subseteq Q$. Well, $\mathbb{R}^n \setminus C$ is open. Thus, $(\mathbb{R}^n \setminus C) \cap Q$ is open in Q . Well, $C = Q \setminus (Q \setminus C)$, so C is closed inside of Q . \square

2.4 Lebesgue’s Number Lemma

30: That is, we are taking a standard cover, not a generalized one.

Let \mathcal{A} be an open cover of some metric space (X, d) so that $\mathcal{A} \subseteq \mathcal{P}(X)$.³⁰ Let $x \in X$. Our goal is to determine if we can cover X in some small balls with equal radius so that each ball is only an $A \in \mathcal{A}$.

Definition 2.4.1 (Lebesgue Number) *The Lebesgue number of the cover \mathcal{A} is any number $\delta > 0$ such that any subset $S \subseteq X$ of $\text{Diam}(S) < \delta$ is contained in A for some $A \in \mathcal{A}$.*

Remark 2.4.1 Looking at the open interval $(0, 1) \subseteq \mathbb{R}$, it is clear that the Lebesgue number does not exist, in general. However, it turns out, compactness gives us exactly what we need.

Definition 2.4.2 (Distance to Set) *For any $C \subseteq X$, define*

$$d(x, C) := \inf_{y \in C} \{d(x, y)\}.$$

Note that there is also the Hausdorff distance which uses supremum.

Theorem 2.4.1 (Lebesgue’s Lemma) *If (X, d) is compact and $\mathcal{A} := \{A_i : i \in I\}$ is an open cover of X , then a Lebesgue number exists.*

31: \mathcal{A}' is a subfamily of \mathcal{A} . Note that if we can find a Lebesgue number for the finite cover, then we get one for \mathcal{A} for free.

32: Since we excluded the empty case, our $d(x, C_k) \neq \infty$.

Proof. There exists a finite subcover $\mathcal{A}' := \{A'_k : 1 \leq k \leq n\}$.³¹ Denote $C_k := X \setminus A'_k$ for any $k \in \{1, \dots, n\}$. Note that if $A'_k = X$, then let $\delta := 1$, or whichever number you like. This is certainly a Lebesgue number for \mathcal{A}' . Assume $C_k \neq \emptyset$ for any k . Consider the function³²

$$\begin{aligned} X &\xrightarrow{f} [0, \infty) \subseteq \mathbb{R} \\ x &\longmapsto \frac{1}{n} \sum_{k=1}^n d(x, C_k). \end{aligned}$$

From the triangle inequality of $d : X^2 \rightarrow [0, \infty)$,

$$d(x, y) - d(x', y) \leq d(x, x')$$

and

$$d(x', y) - d(x, y) \leq d(x, x'),$$

so $|d(x, y) - d(x', y)| \leq d(x, x')$.³³ Similarly, we hope to have

$$|d(x, C) - d(x', C)| \leq d(x, x').$$

We have that, for $c \in C$,

$$d(x, C) \leq d(x, c) \leq d(x, y) + d(y, c)$$

Hence,

$$d(x, C) - d(x, y) \leq d(y, c).$$

The LHS is independent of c , so

$$d(x, C) - d(x, y) \leq \inf_{c \in C} \{d(y, c)\} = d(y, C).$$

Interchanging the roles of x, y we get the opposite statement, so together the absolute value statement holds. Hence, the map

$$d(\cdot, C) : X \rightarrow [0, \infty)$$

is 1-Lipchitz.³⁴ Hence, $f : X \rightarrow [0, \infty)$ is continuous (the sum of finitely many continuous functions to \mathbb{R} is continuous, as proven on the homework), plus $f(x) > 0$, as each A_k is open. Thus, there exists an $x_{\min} \in X$ such that for all $x \in X$, $0 < f(x_{\min}) \leq f(x)$. Denote $\delta := f(x_{\min}) > 0$. Then, for any x , $f(x) \geq \delta$. Let C be any subset of X of diameter less than δ . Let $x_0 \in C$. Then, $C \subseteq B(x, \delta)$. We have that

$$0 < \delta \leq f(x_0) = \frac{1}{n} \sum_{i=1}^n d(x_0, C_i) \leq \max_{1 \leq i \leq n} \{d(x_0, C_i)\}.$$

There exists a number $m \in \{1, \dots, n\}$ so that

$$\max_{1 \leq i \leq n} \{d(x_0, C_i)\} = d(x_0, C_m),$$

meaning $0 < \delta \leq d(x_0, C_m)$. Thus,

$$B(x_0, \delta) \cap C_m = \emptyset \Leftrightarrow B(x_0, \delta) \subseteq A'_m \in \mathcal{A},$$

as desired. Thus, δ is a Lebesgue number for \mathcal{A} . □

Definition 2.4.3 (Uniformly Continuous) *Let $f : X \rightarrow Y$ be a function between metric spaces. We call f uniformly continuous if for all $\varepsilon > 0$ there exists a $\delta > 0$ so that for all $x, x' \in X$,*

$$d_X(x, x') < \delta \implies d_Y(f(x), f(x')) < \varepsilon.$$

It is pretty clear that

$$\text{Uniform Continuity} \implies \text{Continuity}.$$

33: Note that this bound does not depend on y at all. This means $d(\cdot, y) : X \rightarrow [0, \infty)$ is 1-Lipchitz.

34: Note that Lipchitz implies uniform continuity with $\delta := \varepsilon/L$.

Certainly, the converse does not generally hold.

Example 2.4.1 The exponential function \exp on \mathbb{R} is *not* uniformly continuous. It goes wrong, because \mathbb{R} is not compact!

Theorem 2.4.2 *If X is a compact metric space and $f : X \rightarrow Y$ is a continuous function between metric spaces, then f is uniformly continuous.*

Proof. To say that $d_X(x, x') < \delta$, we can say

$$\text{Diam}\{x, x'\} < \delta.$$

Cover Y with open balls of radius $\varepsilon/2 > 0$. By pulling back on f , the set

$$\{f^{-1}(B(y, \varepsilon/2)) : y \in Y\}$$

35: This is where Lebesgue’s lemma is useful.

is an open cover of X . Then, there exists a Lebesgue number δ^{35} for this cover. Fill in the details as an exercise. \square

2.5 Compactifications and Zorn’s Lemma

Definition 2.5.1 (Compactification) *A compactification of a topological space X is any topological space Y so that $X \subseteq Y$ (as a subspace) so that Y is compact, Hausdorff, and $\overline{X}^Y = Y$.*

Definition 2.5.2 (One-Point Compactification) *If $Y \setminus X$, as above, is a singleton, then Y is a one-point compactification.*

Remark 2.5.1 We can compactify an open interval by adding two points and getting the closed interval, or by adding one and getting an object homeomorphic to the circle.

36: This proof was omitted from lecture for brevity.

We briefly give the general outline of proving sequential compactness implies compactness for metrizable (first-countable) spaces.³⁶

- ▶ Prove that sequential compactness implies the existence of a Lebesgue number for any open cover. To do so, proceed by contradiction.
- ▶ Prove that for all $\varepsilon > 0$, there exists a finite cover of X consisting of ε -balls. Again, proceed by contradiction.
- ▶ Combine the previous parts.

Lemma 2.5.1 *Let X be compact and Y be Hausdorff. Define $f : X \hookrightarrow Y$ to be a continuous injection. Restrict the codomain to extract $f' : X \xrightarrow{\text{Set}} f(X)$, a bijection in **Set**. Then, f' is a homeomorphism.³⁷*

37: That is, we can “promote” an isomorphism in **Set** to an isomorphism in **Top**. This is not a rigorous notion, and the actual realizable functor is quite complex.

Proof. Take any closed $C \subseteq X$. Then, $f'(C)$ is compact in $f'(X) \subseteq Y$. Since Y is Hausdorff, then $f'(C)$ is closed in $f(X)$. Thus, $g' = f'^{-1}$ is continuous. \square

We can now give another description of compactness.

Proposition 2.5.2 *The following are equivalent:*

- (i) X is compact; i.e., any open cover of X admits a finite subcover.
- (ii) For any family $\mathcal{C} := \{C_i \subseteq X\}_{i \in I}$ of closed sets so that $\bigcap_I C_i = \emptyset$, there exists a finite subfamily $\{C_{i_1}, \dots, C_{i_n}\}$ so that $\bigcap_n C_{i_j} = \emptyset$.³⁸
- (iii) Let \mathcal{C} be a family of closed subsets in X . If every finite subfamily of \mathcal{C} has nonempty intersection,³⁹ then $\bigcap \mathcal{C} \neq \emptyset$.

38: Duality of complements gives this formulation of compactness clearly.

39: We denote this condition by FIP (finite intersection property). Note that (iii) follows by taking the contrapositive of (ii).

Definition 2.5.3 (Non-Strict Partially Ordered Set) *A (non-strict) partially ordered set (poset) is a pair (S, \leq) so that*

- (i) \leq is reflexive.
- (ii) \leq is antisymmetric.⁴⁰
- (iii) \leq is transitive.

40: If $x \leq y$ and $y \leq x$, then $x = y$.

We call \leq , as above, a non-strict partial order, to be pedantic.⁴¹

41: We will prefer the non-strict convention.

Definition 2.5.4 (Strict Partial Order) *A relation $<$ is a strict partial order if*

- (i) for all $x \in S$, $x \not< x$.
- (ii) for all $x, y \in S$, if $x < y$ then $y \not< x$.
- (iii) for all $x, y, z \in S$, if $x < y$ and $y < z$ then $x < z$.

Remark 2.5.2 The important thing to notice here is that we can interchange between \leq and $<$ freely.

Example 2.5.1 (Posets)

- (i) The class of all topologies \mathcal{T} on a set S , where inclusion \subseteq is the partial order, is a poset.
- (ii) Another standard example is the subgroup lattice, taking all subgroups H of a set G , with a partial order \leq .

Definition 2.5.5 (Total Order) *A total (or simple/linear) order on a set S is a partial order \leq on S if for all $a, b \in S$, $a \leq b$ or $b \leq a$.*

Definition 2.5.6 (Totally Ordered Set) *A totally ordered set (or chain) is a set with a total order (S, \leq) .*

Definition 2.5.7 (Upper Bound) *Given a subset $S' \subseteq S$ where S is a poset, an upper bound on S' in S is an element $u \in S$ so that for all $s' \in S'$, $s' \leq u$.⁴²*

42: Similarly, you could talk about lower bounds.

Definition 2.5.8 (Maximal) *An element $m \in S$ is maximal if for all $s \in S$ so that $m \leq s$, we have $m = s$.⁴³*

43: In totally ordered sets, this is the same as the "largest" element. However, in general they do not coincide. For instance, in \mathbb{R} , every closed subset has a maximal element which is also its largest. Yet, we could have a three-element set $\{a, b, c\}$ so that $a > b$, $c > b$, but a, c are not comparable. Then, both a, c are maximal, but there is no largest.

Lemma 2.5.3 (Zorn's) *Let (S, \leq) be a nonempty poset such that any chain $\mathcal{C} \subseteq S$ has an upper bound in S . Then, S has a maximal element.*

Proof. The lemma is equivalent to the axiom of choice. We will simply assume ZFC hereafter, taking Zorn's for granted. \square

Remark 2.5.3 Let us take a brief aside on notation. Let C be a set. Then, $c \in C$ is an element. Consider now a family $\mathcal{C} \subseteq \mathcal{P}(C)$. In particular, $\mathcal{C} \in \mathcal{P}(\mathcal{P}(C))$. Then, we can form classes/supersets $\mathbf{C} \subseteq \mathcal{P}(\mathcal{P}(C))$, so $\mathbf{C} \in \mathcal{P}(\mathcal{P}(\mathcal{P}(C)))$. This will be useful in our proof of Tychonoff's theorem.

Theorem 2.5.4 (Tychonoff's Theorem) *Let $\{(X_i, \mathcal{T}_i) : i \in I\}$ be a family of compact spaces. Then, the product $X := \prod_I X_i$ with $\mathcal{T}_{\text{prod}}$ is compact.*

Proof. We will use the FIP characterization of compactness. Let \mathcal{A} be a family of closed subsets of X satisfying the FIP. Since every $A \in \mathcal{A}$ is closed, we can write

$$\bigcap_{D \in \mathcal{D}} \overline{D} \subseteq \bigcap_{A \in \mathcal{A}} \overline{A} = \bigcap_{A \in \mathcal{A}} A = \bigcap \mathcal{A},$$

where we hope to construct such a \mathcal{D} , as proving $\bigcap_{\mathcal{D}} D \neq \emptyset$ will show $\bigcap \mathcal{A}$ is nonempty. Let

$$\mathbf{A} := \{\mathcal{A}' \subseteq \mathcal{P}(X) : \mathcal{A} \subseteq \mathcal{A}' \text{ and } \mathcal{A}' \text{ has FIP}\} \subseteq \mathcal{P}(\mathcal{P}(X)).$$

We can place a natural partial order \leq on \mathbf{A} by set theoretic inclusion. We will write $\mathcal{A}' \leq \mathcal{A}''$ if and only $\mathcal{A}' \subseteq \mathcal{A}''$. If $\mathcal{A} = \emptyset$, then $\bigcap \mathcal{A} = X \neq \emptyset$, using the axiom of choice.⁴⁴ Now, we must check that for any chain \mathbf{B} in \mathbf{A} , \mathbf{B} has an upper bound $\mathcal{C} \in \mathbf{A}$. Take any chain $\mathbf{B} \subseteq \mathbf{A}$. Define

$$\mathcal{C} := \bigcup_{\mathcal{B} \in \mathbf{B}} \mathcal{B}.$$

Clearly, for all $\mathcal{B} \in \mathbf{B}$, $\mathcal{B} \subseteq \mathcal{C}$, so \mathcal{C} is an upper bound. We know $\mathcal{A} \subseteq \mathcal{C}$, as $\mathcal{A} \subseteq \mathcal{B}$ for all $\mathcal{B} \in \mathbf{B}$. Take any $C_1, \dots, C_n \in \mathcal{C}$. We want to show

$$C_1 \cap \dots \cap C_n \neq \emptyset.$$

For all $k \in [n]$, there exists a family $\mathcal{B}_k \in \mathbf{B}$ such that $C_k \in \mathcal{B}_k$. Since \mathbf{B} is a chain with respect to \subseteq , there exists a largest family, say $\mathcal{B}_{k'}$.⁴⁵ Thus, $C_1, \dots, C_n \in \mathcal{B}_{k'} \in \mathbf{B} \subseteq \mathbf{A}$. Well, $\mathcal{B}_{k'}$ satisfies the FIP, so $\mathcal{C} \in \mathbf{A}$. By Zorn's lemma, there exists a maximal family $\mathcal{D} \in \mathbf{A}$. In particular, $\mathcal{A} \subseteq \mathcal{D}$, and \mathcal{D} satisfies the FIP. We also have that \mathcal{D} is maximal in

$$\{\mathcal{A}' \subseteq \mathcal{P}(X) : \mathcal{A}' \text{ satisfies FIP}\}.$$

We say that \mathcal{D} is maximal with respect to the FIP. All that remains to be shown is that

$$\bigcap_{D \in \mathcal{D}} \overline{D} \neq \emptyset.$$

44: We select an element in X_i for each index X_i to form $\underline{x} : I \rightarrow \bigcup_I X_i$.

45: That is, for all $k \in [n]$, $\mathcal{B}_k \subseteq \mathcal{B}_{k'}$.

Define the family, for $j \in I$,

$$\{\overline{\pi_j(D)} : D \in \mathcal{D}\} \subseteq \mathcal{P}(X_j).$$

We claim that this family satisfies the FIP. Since X_j is compact,

$$\bigcap_{D \in \mathcal{D}} \overline{\pi_j(D)} \neq \emptyset.$$

Thus, there exists $x_j \in \bigcap_{D \in \mathcal{D}} \overline{\pi_j(D)} \subseteq X_j$. Define $\underline{x}(j) := x_j$.⁴⁶ This gives us a point $\underline{x} \in \prod_I X_i = X$. Pick any $D \in \mathcal{D}$. We want to show $\underline{x} \in \overline{D}$. Take any open neighborhood U_j of x_j . Then, we have the subbasic set $\pi_j^{-1}(U_j)$. We have that $x_j \in U_j \cap \pi_j(D) \neq \emptyset$. Taking preimages, $\pi_j^{-1}(U_j) \cap D \neq \emptyset$. Via maximality and the lemma below, $\pi_j^{-1}(U_j) \in \mathcal{D}$, for any neighborhood U_j of x_j in X_j . Also by the lemma, every basic neighborhood containing \underline{x} in X in $\mathcal{T}_{\text{prod}}$ is in \mathcal{D} . Take any such basic neighborhood U of \underline{x} in X . For all $D \in \mathcal{D}$, we claim that $U \cap D \neq \emptyset$.⁴⁷ Thus, $\underline{x} \in \overline{D}$ for each $D \in \mathcal{D}$, meaning $\underline{x} \in \bigcap_{D \in \mathcal{D}} \overline{D}$, as desired. \square

46: We need choice to turn nonemptiness into a function.

47: Use the FIP for \mathcal{D} .

Lemma 2.5.5 *If $\mathcal{D} \subseteq \mathcal{P}(X)$ is maximal with respect to the FIP, then*

- (i) \mathcal{D} is closed under finite intersections.
- (ii) if $A \subseteq X$ is so that $A \cap D \neq \emptyset$ for all $D \in \mathcal{D}$, then $A \in \mathcal{D}$.

- (i) *Proof.* Pick finitely many $D_1, \dots, D_n \in \mathcal{D}$. We want to show that $D_1 \cap \dots \cap D_n \in \mathcal{D}$. Consider

$$\mathcal{D}' := \mathcal{D} \cup \{D_1 \cap \dots \cap D_n\}.$$

If we can show that $\mathcal{D}' \in \mathbf{A}$, then we win. Certainly $\mathcal{A} \subseteq \mathcal{D}'$. Pick any finite subfamily $D'_1, \dots, D'_m \in \mathcal{D}'$. If $D'_i \in \mathcal{D}$ for all i , then FIP implies $\bigcap_m D'_i \neq \emptyset$. If one of the $D'_m = D_1 \cap \dots \cap D_n$, then

$$D'_1 \cap \dots \cap (D_1 \cap \dots \cap D_n) \neq \emptyset,$$

also by FIP.⁴⁸ Thus, $\mathcal{D}' \in \mathbf{A}$. Yet, $\mathcal{D} \subsetneq \mathcal{D}'$, a contradiction to maximality. \square

48: Taking intersections is associative.

- (ii) *Proof.* Proceed, again, by contradiction to maximality. \square

**ON EMBEDDINGS, FUNCTION SPACES, AND
ALGEBRAIC TOPOLOGY**

Separation and Embeddings

We have previously discussed this notion of being Hausdorff:

Definition 3.0.1 (Axiom T_2 : Hausdorff) *A topological space X is Hausdorff (or T_2) if for all $x \neq y \in X$, there exist neighborhoods U of x and V of y such that $U \cap V = \emptyset$.*

Naturally, we want to both weaken and strengthen this axiom.

3.1 Introducing the T_i Axioms

Definition 3.1.1 (Axiom T_1) *A topological space X satisfies T_1 if for all $x \neq y \in X$, there exists a neighborhood U of x such that $y \notin U$ and there exists a neighborhood V of y such that $x \notin V$.*

Remark 3.1.1 It is clear that $T_2 \Rightarrow T_1$.

Definition 3.1.2 (Axiom T_3 : Regular) *A topological space X is regular (or T_3) if each singleton subset of X is closed, and if for all $x \in X$ and for all closed $C \subseteq X$, if $x \notin C$, then there exists a neighborhood U of x and there exists a neighborhood V of C such that $U \cap V = \emptyset$.¹*

Remark 3.1.2 Since singletons are closed, we have $T_3 \Rightarrow T_2$.

Definition 3.1.3 (Axiom T_4 : Normal) *A topological space X is normal (or T_4) if each singleton subset of X is closed, and for all closed subsets $C_1, C_2 \subseteq X$, if $C_1 \cap C_2 = \emptyset$, then there exist neighborhoods U of C_1 and V of C_2 such that $U \cap V = \emptyset$.*

Remark 3.1.3 Once again, $T_4 \Rightarrow T_3$.

Thus, we have the chain of implications

$$T_4 \implies T_3 \implies T_2 \implies T_1.$$

Definition 3.1.4 (Axiom $T_{3\frac{1}{2}}$: Completely Regular) *A topological space X is completely regular if singletons are closed, and if for any point $x \in X$ and closed subset $C \subseteq X$ such that $x \notin C$, there exists a continuous function $f : X \rightarrow [0, 1]$ such that $f : x \mapsto 0$ and $C \mapsto \{1\}$.²*

We are using that $[0, 1]$ is Hausdorff. Take open intervals around 0 and 1 called V, V' . Then, $f^{-1}(V) \cap f^{-1}(V') = \emptyset$ is an open separation of x and

3.1 Introducing the T_i Axioms . 37

3.2 Embeddings and Stone-Čech 38

1: That is, you can separate any closed set from any point.

2: In some sense, we are separating the point and the set by a function.

the closed set C . As desired, this tells us

$$T_{3\frac{1}{2}} = \text{Completely Regular} \implies T_3.$$

It turns out, that $T_4 \implies T_{3\frac{1}{2}}$, which follows from what is called Urysohn's lemma. We will skip the proof of Urysohn's, though you can find the proof in any standard text.³

3: For instance, see Munkres.

3.2 Embeddings and Stone-Ćech

We now want a way to construct maps which behave like inclusions, with the hope of using these functions to "embed" spaces into Hausdorff compact spaces. As you should remember, if we can force some subspace of our larger space to equal the closure of the embedded space, then we can form a compactification.

Definition 3.2.1 (Embedding) *An embedding of X into a topological space Y is a continuous injection $X \hookrightarrow Y$ such that $f' : X \xrightarrow{\simeq}_{\text{Top}} f(X) \subseteq Y$ is a homeomorphism.*⁴

4: Let $f(X)$ have the subspace topology inherited from Y .

Theorem 3.2.1 (Embedding Theorem) *Let X be a topological space and $\{f_i : X \rightarrow [0, 1]\}_{i \in I}$ be a family of continuous functions such that for all $x \in X$ and neighborhoods U of x , there exists an $i \in I$ so that $f_i(x) > 0$ and $f_i(X \setminus U) = \{0\}$. Then, the function⁵ $F : X \rightarrow [0, 1]^I$ defined by $F(x)(i) = f_i(x)$ is an embedding.*

5: By definition,

$$[0, 1]^I := \prod_{i \in I} [0, 1].$$

Proof. The proof is not particularly illuminating, so check Munkres. \square

Lemma 3.2.2 (Existence of Stone-Ćech) *Let X be completely regular. Then, a compactification exists.*

Proof. Let X be a $T_{3\frac{1}{2}}$ topological space. Let $\{f_i : i \in I\}$ be the set of all continuous bounded functions $f_i : X \rightarrow \mathbb{R}$. Applying the embedding theorem tells us that we have an embedding

$$F : X \rightarrow \prod_{i \in I} [\inf f_i(x), \sup f_i(x)].$$

Now, by the definition of an embedding $X \simeq F(X) \subseteq \overline{F(X)}^\Pi$. Then, $\overline{F(X)}$ is Hausdorff and compact, so it is a compactification of X . \square

Definition 3.2.2 (Stone-Ćech Compactification) *We call $\beta(X) := \overline{F(X)}$, as above, the Stone-Ćech compactification of X .*

Theorem 3.2.3 (Universal Property of Stone-Ćech Compactification) *Let X be completely regular. For all bounded continuous functions $f : X \rightarrow \mathbb{R}$, there exists a continuous function $g : \beta(X) \rightarrow \mathbb{R}$ which extends f via the*

inclusion $X \hookrightarrow \beta(X)$.




Figure 3.1: Universal property of Stone-Čech compactification

Proof. We do not prove the universal property, but Munkres essentially constructs $\beta(X)$ via this property. \square

We claim that if C is compact and Hausdorff, then the Stone-Čech compactification satisfies the diagram






Figure 3.2: Generalized universal property

Again, see Munkres for the proof. Using the universal property as a definition, you can move to arbitrary compact Hausdorff spaces C .⁶

6: We are going to use this for uniqueness, up to homeomorphism.

Theorem 3.2.4 (Uniqueness of Stone-Čech) *Let $\beta_1(X)$ and $\beta_2(X)$ be Stone-Čech compactifications of a $T_{3\frac{1}{2}}$ topological space. Then, $\beta_1 \simeq \beta_2$.*

Proof. Via the generalized universal property, we have the diagrams



Gluing the diagrams together, we get that the diagram

$$\begin{array}{ccccc}
 & & X & & \\
 & \swarrow & \downarrow & \searrow & \\
 \beta_1(X) & \xrightarrow{g_1} & \beta_2(X) & \xrightarrow{g_2} & \beta_1(X)
 \end{array}$$

has $g_2 \circ g_1$ is continuous and equal to id_X . By symmetry, $g_1 \circ g_2$ is continuous and $g_1 \circ g_2 = \text{id}_X$. We want to deduce that $g_2 \circ g_1 = \text{id}_{\beta_1(X)}$ and $g_1 \circ g_2 = \text{id}_{\beta_2(X)}$. We will be done via the following lemma, as $g_1^{-1} = g_2$, so g_1^{-1} is continuous. Thus, g_1 is a homeomorphism.⁷ \square

7: An isomorphism in any category is just an invertible. Recall that in the case of Top, we call these homeomorphisms. This is a homeomorphism extends identity on X .

Lemma 3.2.5 (Uniqueness of Extensions) *Let $A \subseteq X$ and $f : A \rightarrow Y$, where Y is Hausdorff, and $g, g' : \bar{A} \rightarrow Y$ are continuous extensions of f . Then, $g = g'$.*

Proof. Suppose not; i.e., $g \neq g'$. Then, there exists $x \in \bar{A}$ such that $g(x) \neq g'(x)$. Since Y is Hausdorff, we can separate the values by neighborhoods V of $g(x)$ and V' of $g'(x)$ so that $V \cap V' = \emptyset$. By continuity, $U := g^{-1}(V) \cap g'^{-1}(V') =: U'$ is a neighborhood of X . Then, $(U \cap U') \cap A \neq \emptyset$. Pick a point $y \in (U \cap U') \cap A$, then $g(y) \in V$ and $g'(y) \in V'$. Since $y \in A$, $g(y) = f(y) = g'(y)$. Thus, $V \cap V' \neq \emptyset$, a contradiction. \square

Completeness and Function Spaces

4

For this discussion, let (X, d) be a metric space with a corresponding distance function $d : X^2 \rightarrow [0, \infty)$.

4.1 Cauchy Sequences and Completions

Recall a standard definition from analysis.

Definition 4.1.1 (Cauchy Sequence) *A Cauchy sequence in X is a sequence $\{x_n\}$ in X such that for all $\varepsilon > 0$, there exists an $N \in \mathbb{Z}_+$ such that for all $m, n \in \mathbb{Z}_+$, if $n, m \geq N$, then $d(x_n, x_m) < \varepsilon$.*

Proposition 4.1.1 *Any convergent sequence $\{x_n\} \subseteq X$ is Cauchy.*

Example 4.1.1 Consider a sequence in \mathbb{Q} “converging” to $\sqrt{2}$ in \mathbb{R} . This is clearly Cauchy, but not convergent in (\mathbb{Q}, d) .

From this example, it is clear that the converse of the proposition above is not true.

Definition 4.1.2 (Complete) *A metric space (X, d) is complete if any Cauchy sequence in X is convergent in X .*

Our goal is now to construct a larger, complete space to embed a non-complete space within.

Definition 4.1.3 (Completion) *Given a metric space (X, d) , a completion of X is a complete metric space (\hat{X}, \hat{d}) such that $X \subseteq \hat{X}$, $\hat{d} : \hat{X}^2 \rightarrow [0, \infty)$ is a metric extending d along the inclusion $\iota : X \hookrightarrow \hat{X}$, and $\overline{X^{\text{seq}}}^{\hat{d}} = \hat{X}$.¹*

Lemma 4.1.2 *Let (X, d) be any complete metric space. Let Y be a closed subspace $Y \subseteq X$. Then, Y is complete with respect to $d_Y := d|_Y$.*

Proof. Since the metrics agree, we have that $\{y_n\} \subseteq Y$ has $y_n \rightarrow x \in X$. To show $x \in Y$, it suffices to show that $x \in \overline{Y}$. Well, we showed that $\overline{Y^{\text{seq}}} = \overline{Y}$ for metric spaces, so we are done. \square

Lemma 4.1.3 *In the setting above, the converse also holds.*

Proof. Pick $\bar{y} \in \overline{Y}$. Then, pick a sequence $\{y_n\}$ such that $y_n \rightarrow \bar{y} \in \overline{Y}$ and $y_n \rightarrow y' \in Y$. Since Y is Hausdorff, this forces $\bar{y} = y' \in Y$. \square

4.1 Cauchy Sequences and Completions	41
4.2 Bounded Space	42
4.3 More on Function Spaces . .	45
4.4 Arzelà-Ascoli Theorem . . .	48

1: This forces a sort of minimality, like we did with completions.

Lemma 4.1.4 *If $\{x_n\}$ is a Cauchy sequence in a metric space (X, d) , then the following are equivalent:*

- (i) x_n converges in X .
- (ii) x_n has a subsequence x_{n_k} which converges in X .

Proof. The implication (i) \Rightarrow (ii) is trivial. Let $\{x_{n_k}\} \subseteq \{x_n\}$ be a subsequence which converges to x in X . Take any ε . Then, there exists $N \in \mathbb{Z}_+$ such that for all $m, n \geq N$, $d(x_m, x_n) < \varepsilon/2$. Choose $i \in \mathbb{Z}_+$ so that $n_i \geq N$ and $d(x_{n_i}, x) < \varepsilon/2$. Combining these via the triangle inequality, we get that $x_n \rightarrow x$, as desired. \square

4.2 Bounded Space

Let X, Y be metric spaces.

Definition 4.2.1 (Bounded) *A function $f : X \rightarrow Y$ is bounded if $f(X)$ is bounded in Y . We can use that the diameter of $f(X)$ is finite, or that we can enclose it in a finite-radius ball in Y .*

Definition 4.2.2 ($\mathbb{B}(X, Y)$) *We define*

$$\mathbb{B}(X, Y) := \{f : X \rightarrow Y : f \text{ is bounded}\}.$$

Let d be the metric on Y .

Definition 4.2.3 (Uniform Metric) *Given $f, g \in \mathbb{B}(X, Y)$, let*

$$d_\infty(f, g) := \sup_{x \in X} \{d(f(x), g(x))\} < \infty.$$

2: It is finite, as we take all functions in $\mathbb{B}(X, Y)$ to be bounded.

Remark 4.2.1 Pretty easily, d_∞ is a metric on $\mathbb{B}(X, Y)$.²

Our goal is to construct an injective function $\Phi : X \rightarrow \mathbb{B}(X, \mathbb{R})$, sending $a \mapsto (\Phi_a : X \rightarrow \mathbb{R})$. Fix $x_0 \in X$. For all $a \in X$, define $\Phi_a : x \mapsto d(a, x) - d(x_0, x)$. Check that $\Phi_a \in \mathbb{B}(X, \mathbb{R})$. We have that

$$d(a, x) - d(x_0, x) \leq d(a, x_0)$$

and

$$d(x_0, x) - d(a, x) \leq d(a, x_0),$$

so $|\Phi_a(x)| \leq d(a, x_0)$. Thus, $\Phi_a \in \mathbb{B}(X, \mathbb{R})$. Suppose $a \neq b$. Then, $\Phi_a \neq \Phi_b$, we get $\Phi_a(a) \neq \Phi_b(a)$.

3: That is, it is an injection, plus the metric is preserved:

$$d_\infty(\Phi_a, \Phi_b) = d(a, b).$$

Remark 4.2.2 The function $\Phi : a \mapsto \Phi_a$ is an isometric embedding.³

Now, we will work towards proving the following lemma.

Lemma 4.2.1 *Let X be any set and (Y, d) a complete metric space. Then, $(\mathbb{B}(X, Y), d_\infty)$ is complete.*

Let $f : X \rightarrow Y$ be a function. We want to realize f as an element of some Cartesian product. The easiest way to do so, is to consider Y^X . That is,

$$Y^X := \prod_{x \in X} Y \equiv \mathfrak{F}(X, Y).$$

Clearly, $\mathbb{B}(X, Y) \subseteq Y^X$. If we can show completeness of the latter, and then show closedness of the former, we would be done with our lemma. In particular, we will need a metric for our function space.

Definition 4.2.4 (Forced-Boundedness Metric) *Given (Y, d) , define $\bar{d}(y_1, y_2) := \min\{d(y_1, y_2), 1\} \leq 1$.*⁴

4: As an exercise, check that \bar{d} is a metric.

Proposition 4.2.2 *If $a, b, c \in [0, \infty)$. Suppose $a \leq b + c$. Then, $\min\{a, 1\} \leq \min\{b, 1\} + \min\{c, 1\}$.*

Corollary 4.2.3 *The new metric \bar{d} is a bounded metric.*⁵

5: It is trivially bounded by 1.

If $\varepsilon \leq 1$, then

$$\bar{d}(y_1, y_2) \leq \varepsilon \Leftrightarrow d(y_1, y_2) \leq \varepsilon.$$

Corollary 4.2.4 *If Y is complete with respect to a metric d , then it is complete with respect to \bar{d} .*




Figure 4.1: We have two ways of moving from a metric space (Y, d) to a metric space of functions. We can either take the sup or the forced boundedness. It turns out the results are equal. Prove this as an exercise

Definition 4.2.5 (Isometric Embedding) *An isometric embedding is a function $f : X \rightarrow Y$ between metric spaces such that for all $x, x' \in X$,*

$$d_Y(f(x), f(x')) = d_X(x, x').$$

This implies that X is injective.

Now, we have a method of constructing a completion via

$$\begin{aligned} X &\hookrightarrow \mathbb{B}(X, \mathbb{R}) \\ a &\longmapsto \Phi_a. \end{aligned}$$

Lemma 4.2.5 *If (Y, d) is a complete metric space, then Y^X is complete with respect to \bar{d}_∞ .*

Proof. Let $\{f_n\}$ be any Cauchy sequence in (Y^X, \bar{d}_∞) . For all $\varepsilon > 0$, there exists $N \in \mathbb{Z}_+$ so that for all $m, n \geq N$,

$$\bar{d}_\infty(f_m, f_n) < \varepsilon.$$

Then, for all $x \in X$,

$$\bar{d}(f_m(x), f_n(x)) \leq \bar{d}_\infty(f_m, f_n) < \varepsilon.$$

Thus, for each $x \in X$, the sequence $\{f_n(x)\} \subseteq Y$ is Cauchy. Hence, $\{f_n(x)\}$ converges in Y^6 to a point y_x . Define the function $f : X \rightarrow Y$ be defined by $x \mapsto y_x$. We need to check that $f_n \rightarrow f$ with respect to \bar{d}_∞ . Take any $\varepsilon > 0$. There exists $N \in \mathbb{Z}_+$ such that for all $m, n \geq N$,

$$\bar{d}_\infty(f_m, f_n) < \frac{\varepsilon}{2}.$$

Again, for all $x \in X$,

$$\bar{d}(f_m(x), f_n(x)) \leq \bar{d}_\infty(f_m, f_n) < \frac{\varepsilon}{2},$$

and taking $m \rightarrow \infty$, we get

$$\bar{d}(f_m(x), f_n(x)) \xrightarrow{m \rightarrow \infty} \bar{d}(y_x, f_n(x)) = \bar{d}(f(x), f_n(x)),$$

as $\bar{d} : Y^2 \rightarrow [0, \infty)$ is continuous in the first variable. Then,

$$\bar{d}(f(x), f_n(x)) < \frac{\varepsilon}{2}$$

for all $x \in X$. Thus,⁷

$$\bar{d}_\infty(f, f_n) \leq \frac{\varepsilon}{2} < \varepsilon.$$

□

6: Since d and \bar{d} agree for small values, completeness in (Y, d) implies completeness in (Y, \bar{d}) .

7: Taking the supremum gives us the non-strict inequality, and then we get our strict inequality trivially.

Definition 4.2.6 (Continuous Space) *Let X be a topological space and Y be a topological space. Then, $\mathcal{C}(X, Y)$ is the set of continuous functions $f : X \rightarrow Y$.*

Theorem 4.2.6 *Let X be a topological space and Y be a metric space. Then, both $\mathcal{C}(X, Y)$ and $\mathbb{B}(X, Y)$ are closed in (Y^X, \bar{d}_∞) .⁸*

8: We also want to show that $\mathbb{B}(X, Y)$ is bounded with respect to d_∞ .

Proof. We first show that $\mathcal{C}(X, Y)$ is sequentially closed in Y^X . Suppose $\{f_n\} \subseteq \mathcal{C}(X, Y)$ is a sequence that converges to $f \in Y^X$. We want to use the uniform limit theorem. For all $\varepsilon > 0$, there exists an $n \in \mathbb{Z}_+$ such that for all $n \geq N$,

$$\bar{d}_\infty(f_n, f) < \varepsilon.$$

Again, for all $x \in X$,

$$\bar{d}(f_n(x), f(x)) \leq \bar{d}_\infty(f_n, f) < \varepsilon,$$

so $f_n \rightarrow f$ uniformly. We have shown, $\mathcal{C}(X, Y)$ is sequentially closed, and thus, is closed. Now, let $\{f_n\} \subseteq \mathbb{B}(X, Y)$ be a sequence such that $f_n \rightarrow f \in Y^X$. We want to show $f \in \mathbb{B}(X, Y)$. Let $\varepsilon = 1$. Then, there exists $N \in \mathbb{Z}_+$ such that for all $m, n \geq N$, $d_\infty(f_m, f_n) < 1$. Then, as before,

$$d(f_m(x), f_n(x)) < 1,$$

so take the limit $m \rightarrow \infty$ and get

$$d(f(x), f_n(x)) \leq 1.$$

We know f_n is bounded. Thus, f is bounded. \square

Remark 4.2.3 Inspired by the low-dimensional topology *Coble Lectures* in Altgeld Hall, we will briefly mention n -manifolds.⁹

9: The lectures are presented by Peter Kronheimer. He will focus on 3- and 4-manifolds.

Definition 4.2.7 (n -Manifold) *An n -manifold is a topological space M such that*

- (i) M is locally Euclidean.¹⁰
- (ii) M is Hausdorff (T_2) and second-countable.

10: That is, for all $x \in M$, there exists a neighborhood U of x in M such that U is homeomorphic to $\mathbb{B}^n \simeq \mathbb{R}^n$.

Example 4.2.1 Consider the image of the embedding $\mathbb{S}^1 \hookrightarrow K$, the right-trefoil knot. Then, $\mathbb{R}^3 \setminus K$ is a 3-manifold.

Proposition 4.2.7 *Every compact metric space is complete.*

Proof. Take any Cauchy sequence. Since the space is compact and metrizable, it is sequentially compact, so the sequence has a convergent subsequence. We showed this is equivalent to the sequence converging, so the space is complete. \square

4.3 More on Function Spaces

If X is a set and Y is a topological space, denote for $x \in X$ and an open set $U \subseteq Y$,

$$S(x, U) := \{f : X \rightarrow Y : f(x) \in U\} \subseteq Y^X.$$

Well, for any $f \in Y^X$, if we take $x \in X$, then $f(x) \in U := Y$, so $S(x, U)$ is a subspace.

Definition 4.3.1 (Topology of Pointwise Convergence) *Define a subspace*

$$\mathcal{S}_{point} := \{S(x, U) : x \in X, U \in \mathcal{T}_Y\}.$$

Then, take all finite intersections to get \mathcal{B}_{point} . Finally, take all unions to get

\mathcal{T}_{point} , the topology of pointwise convergence.

Now, let X be a set and Y a metric space. Let $C \subseteq X$

Definition 4.3.2 (Topology of Compact Convergence) *Define*

$$B_C(f, \varepsilon) = \{g : X \rightarrow Y : \sup\{d(f(x), g(x)) : x \in C\} < \varepsilon\}.$$

Then, let $\mathcal{B}_{comp\ conv}$ be the basis of all such $B_C(f, \varepsilon)$. Then, $\mathcal{T}(\mathcal{B}_{comp\ conv}) =: \mathcal{T}_{comp\ conv}$ is the topology of compact convergence.

We have \mathcal{T}_{unif} when our basis elements are $B_X(f, \varepsilon)$, ranging over the whole space.

Theorem 4.3.1 *Let X be a space. Let (Y, d) be a metric space. For the function space Y^X , one has the following inclusion of topologies:¹¹*

$$\mathcal{T}_{unif} \supseteq \mathcal{T}_{comp\ conv} \supseteq \mathcal{T}_{point}.$$

11: This was proven on a day when I slept-in through lecture at 3:00 PM. Check Munkres for his proof.

Remark 4.3.1 If X is compact, then $\mathcal{T}_{unif} = \mathcal{T}_{comp\ conv}$. If X is discrete, then $\mathcal{T}_{comp\ conv} = \mathcal{T}_{point}$.¹²

12: Thus, we have a relation between all three of our new topologies on Y^X . We shall now construct a fourth.

Definition 4.3.3 (Compact-Open Topology) *Define*

$$S(C, U) := \{f \in \mathcal{C}(X, Y) : f(C) \subseteq U\}.$$

Let $\mathcal{S}_{comp\ open}$ be the subbasis of all such elements, where C is compact in X and U is open in Y .¹³ As usual, take finite intersections and arbitrary unions to get $\mathcal{T}_{comp\ open}$, either on Y^X or $\mathcal{C}(X, Y)$.

13: We could just as well look at all of Y^X . Instead we mostly follow Munkres.

Consider $\mathcal{T}_{comp\ open}$ on $\mathcal{C}(X, Y)$, where X is a topological space and Y is a metric space. Our goal is to show that $\mathcal{T}_{comp\ conv}$ on Y^X , restricted to the subspace $\mathcal{C}(X, Y)$, coincides with $\mathcal{T}_{comp\ open}$.

Theorem 4.3.2 *If X is a topological space and (Y, d) is a metric space, then $\mathcal{T}_{comp\ conv}$ and $\mathcal{T}_{comp\ open}$ agree on $\mathcal{C}(X, Y) \subseteq Y^X$.*

Proof. We are looking at the subbasic elements $S(C, U)$ and basic elements $B_C(f, \varepsilon) \cap \mathcal{C}(X, Y)$. We want to show that $\mathcal{T}_{comp\ conv} \subseteq \mathcal{T}_{comp\ open}$. Take any $B_C(f', \varepsilon')$ for some choice of C, f', ε' . Let $f \in B_C(f', \varepsilon')$. There exists a $\varepsilon > 0$ such that $f \in B_C(f, \varepsilon) \subseteq B_C(f', \varepsilon')$.¹⁴ We want to choose C_1, \dots, C_n and U_1, \dots, U_n so that

14: Namely, take

$$\varepsilon := \varepsilon' - \sup_{x \in C} \{d(f(x), f'(x))\}.$$

$$f \in S(C_1, U_1) \cap \dots \cap S(C_n, U_n) \subseteq B_C(f, \varepsilon).$$

For each $x \in C$, choose a neighborhood V_x of x such that $\text{Diam}(f(\overline{V_x})) < \varepsilon$. To find this, first consider $U_x := B(f(x), \varepsilon/4) \subseteq Y$. Then, there exists a neighborhood V_x such that $f(V_x) \subseteq U_x$. Then, $f(\overline{V_x}) \subseteq \overline{B(f(x), \varepsilon/4)}$, by

continuity. Well, $\overline{B(f(x), \varepsilon/4)} \subseteq B(f(x), \varepsilon/3)$. Thus,

$$\text{Diam}(f(\overline{V_x})) \leq \text{Diam}(B(f(x), \varepsilon/3)) \leq \frac{2\varepsilon}{3} < \varepsilon.$$

Since C is compact, there exists a finite subcover $\{V_{x_1}, \dots, V_{x_n}\}$ of C . Denote $C_i := C \cap \overline{V_{x_i}}$, where $i \in \{1, \dots, n\}$. Then, C_i is compact. We have that $C = \bigcup_n C_i$. Also, $\text{Diam}(f(C_i)) \leq \text{Diam}(f(\overline{V_{x_i}})) < \varepsilon$. Let $U_i := U_{x_i} = B(f(x_i), \varepsilon/4)$. We claim that

$$f \in \bigcap_{i=1}^n S(C_i, U_i) \subseteq B_C(f, \varepsilon).$$

The fact that f is in all the $S(C_i, U_i)$ is true by construction.¹⁵ Take any $g \in \bigcap_n S(C_i, U_i)$. We want to show $g \in B_C(f, \varepsilon)$.¹⁶ We have that $g(x), f(x) \in U_{x_i}$, so $d(g(x), f(x)) < \varepsilon$. Since the C_i s cover C , for any $x \in C$, $d(f(x), g(x)) < \varepsilon$, meaning $g \in B_C(f, \varepsilon)$. We now proceed for the converse inclusion. Take any $S(C, U)$, where C is compact in X and U is open in Y . Also, take any $f \in S(C, U)$. Our goal is to find $B_C(f, \varepsilon)$. In particular, we will take f and C to be the same, only searching for ε . Consider $d(f(x), Y \setminus U) := \inf\{d(f(x), y) : y \in Y \setminus C\}$. Furthermore, we claim that $d(\cdot, Y \setminus U)$ is continuous in y' , as d is 1-Lipchitz. Then, $d(\cdot, Y \setminus U) \circ f$ is continuous, and is a function on C . Thus, there exists a minimum; i.e., there exists $x_{\min} \in C$ such that for all $x \in C$,¹⁷

$$0 < d(f(x_{\min}), Y \setminus U) \leq d(f(x), Y \setminus U).$$

Define $\varepsilon := d(f(x_{\min}), Y \setminus U)$. Then,

$$f \in B_C(f, \varepsilon) \cap \mathcal{C}(X, Y) \subseteq S(C, U),$$

meaning $\mathcal{T}_{\text{comp open}} = \mathcal{T}_{\text{comp conv}}$ on $\mathcal{C}(X, Y)$. □

Corollary 4.3.3 *If X and Y are spaces, and $d : X^2 \rightarrow [0, \infty)$ is any metric inducing the topology on Y , then $\mathcal{T}_{\text{comp conv}}$ on $\mathcal{C}(X, Y)$ is independent of d .*

Corollary 4.3.4 *If X is compact, then $\mathcal{T}_{\text{unif}}$ on $\mathcal{C}(X, Y)$ is the same as $\mathcal{T}_{\text{comp open}}$.*

Consider sets X, Y, Z . Construct $Y^{X \times Z}$ of functions $f : X \times Z \rightarrow Y$. On the other hand, consider $(Y^X)^Z$, which is all functions $F : Z \rightarrow Y^X$. As sets, we have a bijection

$$\begin{aligned} Y^{X \times Z} &\xrightarrow[\text{bijection}]{\Phi} (Y^X)^Z \\ f &\longmapsto F(z)(x) := f(x, z), \end{aligned}$$

so $\Phi(f) := F$. Now, let us move to Top:

$$\mathcal{C}(X \times Z, Y) \xrightarrow{\Phi} \mathcal{C}(Z, \mathcal{C}(X, Y)).$$

Put $\mathcal{T}_{\text{comp open}}$ on $\mathcal{C}(X, Y)$. We want to show that $\Phi : f \mapsto F$ is bijective if X is locally compact and Hausdorff.

15: We have $f : V_x \mapsto U_x$ via inclusion, so $f : C_i \mapsto U_i$ via inclusion.

16: We will check that for all $x \in C_i$, $d(g(x), f(x)) < \varepsilon$.

17: It is strictly bounded by zero, using sequential closedness.

18: That is, we can continuously “deform” f into g via h .

Definition 4.3.4 (Homotopy) *Any element $h \in \mathcal{C}(X \times [0, 1], Y)$ is called a homotopy of $f, g \in \mathcal{C}(X, Y)$ if $h(x, 0) = f(x)$ and $h(x, 1) = g(x)$.*¹⁸

Using the above bijection

$$\mathcal{C}(X \times [0, 1], Y) \longrightarrow \mathcal{C}([0, 1], \mathcal{C}(X, Y)),$$

any homotopy can be viewed as a family of functions parameterized by the unit interval $[0, 1]$.

4.4 Arzelà-Ascoli Theorem

We now state a neat theorem which can be applied nicely to the theory of manifolds. Let X be a topological space and Y be a metric space.

Definition 4.4.1 (Surface) *A surface is a 2-manifold.*

19: For \mathcal{F} to be equicontinuous everywhere, the statement holds for all $x \in X$,

Definition 4.4.2 (Equicontinuous) *We say that $\mathcal{F} \subseteq \mathcal{C}(X, Y)$ is equicontinuous at $x_0 \in X$ if for all $\varepsilon > 0$ there exists a neighborhood U of x_0 such that for all $f \in \mathcal{F}$,*¹⁹

$$f(U) \subseteq B(f(x_0), \varepsilon).$$

Definition 4.4.3 (Relatively Compact) *If $A \subseteq Y$, we call A relatively compact if \overline{A} is compact in Y .*

Put the topology of compact convergence on $\mathcal{C}(X, Y)$. Remember, this is the same as the compact-open topology. Note that in \mathbb{R}^n , boundedness is very close to being relative compactness, so the Arzelà-Ascoli theorem is stated using this notion for $\mathcal{C}(X, \mathbb{R}^n)$. Now, let $\mathcal{F} \subseteq \mathcal{C}(X, Y)$. Pick $x_0 \in X$. Then, consider the set

$$S := \{f(x_0) : f \in \mathcal{F}\} \subseteq Y.$$

We say that \mathcal{F} is *pointwise relatively compact* if S is relatively compact for any choice of x_0 .

20: The converse holds if X is locally compact Hausdorff.

Theorem 4.4.1 (Arzelà-Ascoli I) *If $\mathcal{F} \subseteq \mathcal{C}(X, Y)$ is equicontinuous and pointwise relatively compact, then it is relatively compact.*²⁰

21: We say X is σ -compact if it is the countable union of compact spaces. Alternatively, per Munkres, X is Hausdorff σ -compact if there exists a sequence of open subsets U_n and a sequence of compact subspaces C_n such that $U_n \subseteq C_n$. Plus, $\bigcup U_n = X$.

Theorem 4.4.2 (Arzelà-Ascoli II) *Let X be Hausdorff and σ -compact.²¹ Let f_n be a sequence in $\mathcal{C}(X, Y)$ which is equicontinuous and pointwise relatively compact. Then, there is a subsequence of f_n in $\mathcal{C}(X, Y)$.*

We have that I \Rightarrow II.

Introduction to Algebraic Topology

5

Consider the square $[0, 1]^2 \subseteq \mathbb{R}^2$ with the subspace topology. Label opposite sides of the square by a, b respectively, orienting the b -sides up and the a -sides to the right. Then, define an equivalence relation \sim where b -sides are equivalent, a -sides are equivalent, the corners are equivalent, and no points on the interior are equivalent. Taking the quotient space $[0, 1]^2 / \sim$ with the standard quotient topology, we get what is called the 2-torus \mathbb{T}^2 .

On the other hand, we could consider the set $\mathbb{S}^1 \times \mathbb{S}^1$ with each half of the product inheriting their respective \mathbb{R} -subspace topologies. Then, putting the product topology on $\mathbb{S}^1 \times \mathbb{S}^1$, this is *also* \mathbb{T}^2 .

Consider a group action of \mathbb{Z}^2 on \mathbb{R}^2 by

$$\begin{aligned} \mathbb{Z}^2 \times \mathbb{R}^2 &\xrightarrow{\text{action } +} \mathbb{R}^2 \\ (c, d) \cdot (x, y) &\longmapsto (x + c, y + d). \end{aligned}$$

The integer lattice of $[(x, y)]$ is precisely the *orbit* of the additive action. Then, take the quotient $\mathbb{R}^2 / \mathbb{Z}^2$, with cosets given by the equivalence relation of the orbit of the action.¹ This is \mathbb{T}^2 .

Finally, consider the rotation of the circle $\mathbb{S}^1 \subseteq \mathbb{R}^3$ around the z -axis. This is also the same 2-torus \mathbb{T}^2 .

Theorem 5.0.1 *The four definitions of \mathbb{T}^2 , $[0, 1]^2 / \sim$, $\mathbb{S}^1 \times \mathbb{S}^1$, $\mathbb{R}^2 / \mathbb{Z}^2$, and as a solid of revolution are homeomorphic.*

The equivalence of multiple definitions of \mathbb{T}^2 motivates our discussion of algebraic topology. Broadly speaking, algebraic topology precisely studies what its name suggests: how to take algebraic structures like groups, rings, modules, fields, and associate topological spaces with them. We begin this study with the *fundamental group*.

Remark 5.0.1 (Cell Complex on \mathbb{T}^2) Another way to construct \mathbb{T}^2 is as a cell complex. Take one copy of \mathbb{D}^0 , two copies of \mathbb{D}^1 , and one copy of \mathbb{D}^2 . Glue $\partial\mathbb{D}^1$ of both copies to the point, yielding a shape resembling ∞ . Then, glue the a -sides and the b -sides of \mathbb{D}^2 to half of each side of ∞ .²

Similarly, we could form the Δ -complex structure on \mathbb{T}^2 , taking one vertex, three 2-simplices, and two 3-simplices.³ The issue with Δ -complexes is that we lose our information about our 2-simplices. The nice thing about *simplicial-complexes* is that we keep all information about the respective vertices, mapping the simplices injectively (any two points in the simplex should not be identified as one). One way to form the simplicial complex of \mathbb{T}^2 is to start with 4 vertices, 12 edges, and 8 triangles. Prescribe all the gluings as you would expect.

5.1 Fundamental Group 50

5.2 Covering Spaces 53

1: Recall that orbits are just the equivalence classes of the action.

2: This is essentially the same as $[0, 1]^2 / \sim$, but we wanted emphasize the cell structure.

3: If a 3-simplex is a triangle, then a 4-simplex is a tetrahedron (the convex hull of four points in \mathbb{R}^3).

Definition 5.0.1 (Euler Characteristic) *Assigning alternating parity to the n -simplices of the complexes, starting with $+$ on the vertices, the Euler characteristic χ takes the number of each and sums via the parity.*

4: It takes some fair amount of work to prove this generally.

Theorem 5.0.2 *The Euler characteristic of \mathbb{T}^2 is $\chi = 0$.⁴*

It takes some effort, but proving the Euler characteristic is an invariant shows that $\mathbb{T}^2 \not\cong \mathbb{S}^2$.

5.1 Fundamental Group

Let X be a topological space with a base point $x_0 \in X$.

Definition 5.1.1 (Loop) *A loop at $x_0 \in X$ is a path $f : [0, 1] \rightarrow X$ such that $f(0) = x_0$ and $f(1) = x_0$.*

We consider the set $\text{Loops}(X, x_0) \subseteq \text{Paths}(X)$.

5: For the former, we need the starting point of g to be the end point of f .

Definition 5.1.2 (Concatenation Operation) *Let $f, g \in \text{Paths}(X)$ or in $\text{Loops}(X, x_0)$.⁵ Suppose $f(1) = g(0)$. Define a new path $f * g$ by: divide $[0, 1]$ into $[0, 1/2]$ and $(1/2, 1]$. Then consider the linear maps $[0, 1/2] \rightarrow [0, 1]$ prescribed by $t \mapsto 2t$ and $(1/2, 1] \rightarrow [0, 1]$ prescribed by $t \mapsto 2t - 1$. Then, composing, our formula is⁶*

6: This is well-defined, since we took that $f(2(1/2)) = f(1) = g(0) = g(2(1/2) - 1)$.

$$(f * g)(t) := \begin{cases} f(2t), & t \in [0, 1/2] \\ g(2t - 1), & t \in [1/2, 1]. \end{cases}$$

7: It breaks even at associativity.

We claim that $(\text{Loops}(X, x_0), *)$, where $*$ is the concatenation binary operation, is *not* a group.⁷ It looks like we need to be a bit more careful. Given continuous maps $\varphi, \psi : X \rightarrow Y$, recall that a homotopy between φ and ψ is the continuous map

$$H : X \times [0, 1] \rightarrow Y$$

such that $H(x, 0) = \varphi(x)$ and $H(x, 1) = \psi(x)$ for all $x \in X$.

8: We use s for the parameter of the path and t for the parameter of the homotopy.

Definition 5.1.3 (Path Homotopy) *When X happens to also be an interval $X := [0, 1]$, we can consider two paths $f, g : [0, 1] \rightarrow Y$.⁸ A path homotopy from f to g is a continuous function*

$$H : [0, 1] \times [0, 1] \rightarrow Y$$

such that for all $s \in [0, 1]$, $H(s, 0) = f(s)$ and $H(s, 1) = g(s)$. Additionally, we require that for any $t \in [0, 1]$, $H(0, t) = f(0)$ and $H(1, t) = f(1)$.⁹

9: That is, path homotopies preserve the end points.

Definition 5.1.4 (Homotopic) *Two paths $f, g : [0, 1] \rightarrow Y$ are (path-) homotopic if there exists a path homotopy from f to g . We will write $f \sim g$.*

Example 5.1.1 (Trivial Loop) Consider (\mathbb{S}^1, x_0) . One loop is $c_0 : [0, 1] \rightarrow \mathbb{S}^1$, the constant map to x_0 .

Theorem 5.1.1 *Being homotopic is an equivalence relation on $\text{Loops}(X, x_0)$.*¹⁰

10: Or, take $\text{Paths}(X)$.

Proof.

- (i) *Reflexive:* Take $f \in \text{Paths}(X)$. We have the path homotopy $H(s, t) := f(s)$. This is continuous, because it is the composition of the projection from the square to the interval and f .
- (ii) *Symmetry:* Suppose $f \sim g$. Then, there exists a path homotopy H from f to g . Let H' be defined by $H'(s, t) := H(s, 1 - t)$.¹¹
- (iii) *Transitive:* Suppose $f \sim g$ and $g \sim h$. There exists a path homotopy H_1 from f to g and H_2 from g to h . Define

11: Our H' is a composition $(s, t) \mapsto (s, 1 - t) \mapsto H(s, 1 - t)$. The end point mapping property is trivially true.

$$H := \begin{cases} H_1(s, 2t), & t \in [0, 1/2] \\ H_2(s, 2t - 1), & t \in [1/2, 1], \end{cases}$$

noting that at $1/2$, $H(s, 1/2) = H_1(s, 1) = H_2(s, 0) = g(s)$.

□

Definition 5.1.5 (Fundamental Group) *Define the fundamental group $\pi_1(X, x_0) := \text{Loops}(X, x_0)/\sim$, where \sim is homotopy equivalence.¹² The operation $\cdot : \pi_1(X, x_0)^2 \rightarrow \pi_1(X, x_0)$ is defined by $[f] \cdot [g] = [f * g]$.*

12: That is, an element $[f]$ in $\pi_1(X, x_0)$ is a path homotopy equivalence class of loops.

Proposition 5.1.2 *The operation \cdot on $\pi_1(X, x_0)$ is well-defined.*

Proof. Suppose $f' \sim f$ and $g' \sim g$. Then, there exists a path homotopy H_1 between f' and f , and likewise with H_2 and g, g' . Define H by crushing together the H_i horizontally in s , rather than in t :

$$H(s, t) := \begin{cases} H_1(2s, t), & s \in [0, 1/2] \\ H_2(2s - 1, t), & s \in [1/2, 1]. \end{cases}$$

Thus, $[f * g] = [f' * g']$, meaning the operation is well-defined. □

Lemma 5.1.3 *The pair $(\pi_1(X, x_0), \cdot)$ is a group.*

Proof.

- (i) *Associative:* We need to check that

$$([f] \cdot [g]) \cdot [h] = [f] \cdot ([g] \cdot [h]).$$

Well, $([f] \cdot [g]) \cdot [h] = [(f * g) * h]$. Similarly, $[f * (g * h)]$.¹³

13: Again, it will suffice to show that $f * (g * h)$ is path homotopic to $(f * g) * h$. We omit the proof. Define a path homotopy for the intervals $s \in [0, (t + 1)/4]$, $s \in [(t + 1)/4, (t + 2)/4]$, and $s \in [(t + 2)/4, 1]$.

(ii) *Identity*: Our candidate is $c_0 : [0, 1] \rightarrow X$ defined by $[0, 1] \mapsto \{x_0\}$. After concatenation, this is not *exactly* the same, but our goal is to show via path homotopy that

$$[f] \cdot [c_0] = [f * c_0] = [f],$$

meaning $f * c_0 \sim f$. Define the homotopy

$$H(s, t) := \begin{cases} f(\text{a linear map to } [0, 1]), & s \in [0, (t + 1)/2] \\ x_0, & s \in [(t + 1)/2, 1]. \end{cases}$$

This is a path homotopy, so we have the identity $[c_0] \in \pi_1(X, x_0)$.

(iii) *Inverses*: Let $[f] \in \pi_1(X, x_0)$. Let $\bar{f} : [0, 1] \rightarrow X$ be a loop at x_0 defined by $\bar{f}(s) := f(1 - s)$.¹⁴ Again, you can write the formula for $H(s, t)$, defined on the intervals $s \in [0, (1 - t)/2]$, $s \in [(1 - t)/2, (t + 1)/2]$, and $s \in [(t + 1)/2, 1]$. Thus, $[f] \cdot [\bar{f}] = [c_0]$. Then, just interchange the roles of f, \bar{f} , getting $[\bar{f}] \cdot [f] = [c_0]$.

14: This is just us flipping the interval, so we can trace backwards along the path.

Thus, $\pi_1(X, x_0)$ is a group under \cdot . □

Remark 5.1.1 Note that our fundamental group requires a base point. Thus, $\pi_1 : \text{Top}_* \rightarrow \text{Grp}$ is a functor, rather than from Top .

Theorem 5.1.4 $\pi_1(\mathbb{R}^2, x_0) \simeq \{e\}$, the trivial group.

Proof. Let $[f] \in \pi_1(\mathbb{R}^2, x_0)$. Define

$$\ell(t) := (1 - t)f(s) + tx_0.$$

15: It should be clear that this is continuous.

Then, we get a path homotopy¹⁵

$$H(s, t) := (1 - t)f(s) + tx_0.$$

□

Theorem 5.1.5 $\pi_1(\mathbb{T}^2, x_0) \simeq C_\infty \times C_\infty$.

Proof. We will do this informally. Recall the two paths a, b on the square forming \mathbb{T}^2 . These paths satisfy $[a, b] = aba^{-1}b^{-1} = 1$, so we get¹⁶

$$\pi_1(\mathbb{T}^2, x_0) \simeq \langle a, b | [a, b] \sim c_0 \rangle \simeq C_\infty \times C_\infty \simeq \mathbb{Z}^2.$$

16: The two path generators a, b individually generate the free group, but we have to quotient by the relation of the commutator.

□

Example 5.1.2 A few more isomorphisms of fundamental groups:

- (i) $\pi_1(\mathbb{T}^1, x_0) \simeq C_\infty$.
- (ii) $\pi_1(\mathbb{S}^2, x_0) \simeq \{e\}$.
- (iii) $\pi_1(R, x_0) \simeq F(2)$, the free group on two generators, where R is the two-rose.

5.2 Covering Spaces

What is $\pi_1(\mathbb{S}^1, x_0)$. We claim it is $C_\infty \simeq \mathbb{Z}$. The identity, per usual, is the constant loop $[c_0]$. Then, t would be one traversal around \mathbb{S}^1 in one direction. The group is generated by t , with negatives coming from traversing in the opposite direction. How do we do show this?¹⁷

17: We will sketch out the basic idea.

Well, consider $\mathbb{R} \rightarrow \mathbb{S}^1 \subseteq \mathbb{C}$. Then, we write $f : [0, 1] \rightarrow \mathbb{S}^1 \subseteq \mathbb{C}$, sending $s \mapsto \cos(2\pi s), \sin(2\pi s)$. Then, $f^5 = f * \dots * f$ is given by $s \mapsto (\cos 10\pi s, \sin 10\pi s)$. The question is why if we take any loop, then it will be path homotopic to one of this flavor. For any small open interval on \mathbb{R} , you can see that we would have a corresponding small ball on \mathbb{S}^1 . Then, slowly traversing our path, we can lift each coordinate up into \mathbb{R} . Using this lifting (lift x_0 to 0) and compactness, the other endpoint will be at $-n$, where we go around n times. That is, $\mathbb{Z} \subseteq \mathbb{R}$ is the preimage of x_0 . The map from the line to the circle is an important example of a covering space, lifting paths piece by piece.

Informally, a *covering space* of X is a function $p : \tilde{X} \rightarrow X$ which is a local homeomorphism.¹⁸

18: This is *not* equivalent to the actual definition. We are just looking for intuition.

Remark 5.2.1 For a particular type of covering spaces, called regular covers, the $\pi_1(X, x_0) =: G$ acts on the covering space, in the way \mathbb{Z} acts on \mathbb{R} by translation.

Definition 5.2.1 (Covering Space) A covering $p : \tilde{X} \rightarrow X$ of X is a continuous map such that for all $x \in X$, there exists a neighborhood U_x of x and a discrete topological space D_x so that the preimage $p^{-1}(U_x) = \coprod_{D_x} V_k$, where $p|_{V_k} : V_k \rightarrow U_x$ is a homeomorphism for all k .

Consider a circle which has 1/2 the speed of traversal as \mathbb{S}^1 . This is *also* a covering space of the circle \mathbb{S}^1 , under the projection map $\times 2$. That is, antipodal points correspond to a single point on \mathbb{S}^1 , precisely like our definition of \mathbb{RP}^2 . We could continue this process for all $\times n$ maps for $n \in \mathbb{Z}$. Then, we get corresponding subgroups associated with $n\mathbb{Z}$.

Definition 5.2.2 (Simply Connected) A space X is simply connected if X is path connected and $\pi_1(X) = 1$.

Among all covering spaces for \mathbb{S}^1 , the only simply connected one is \mathbb{R} . In this case, we call \mathbb{R} the *universal covering space* of \mathbb{S}^1 . What is a universal covering space of \mathbb{RP}^2 ?¹⁹ It turns out, the answer is precisely \mathbb{S}^2 , as one would expect.

19: In fact, universal covering spaces are unique up to homeomorphism.

What is the covering space of the 2-rose? We know the fundamental group is $F(2)$, the free group on two generators. If we trace up the fibers to a covering space, we could form a space which has an infinite amount of 4-branches. Then, we could either wrap around or make a *tree*. Recall that any subgroup of the free group is another free group.²⁰

20: That is, subgroups of the fundamental group correspond to covering spaces.